

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
14 March 2002 (14.03.2002)

PCT

(10) International Publication Number
WO 02/20489 A2

- (51) International Patent Classification: **C07D 215/00**
- (21) International Application Number: PCT/US01/26130
- (22) International Filing Date: 21 August 2001 (21.08.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/230,267 6 September 2000 (06.09.2000) US
- (71) Applicant (for all designated States except US): **BRISTOL-MYERS SQUIBB COMPANY** [US/US]; P.O. Box 4000, Lawrenceville-Princeton Rd., Princeton, NJ 08543-4000 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **BI, Yingzhi** [CN/US]; 5 Mershon Lane, Plainsboro, NJ 08536 (US). **YU, Guixue** [US/US]; 8 Greene Drive, Lawrenceville, NJ 08548 (US). **ROTELLA, David, P.** [US/US]; 1360 Fountain Road, Newton, PA 18940 (US). **MACOR, John, E.** [US/US]; 644 Moose Hill Rd., Guilford, CT 06437 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EL, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 02/20489 A2

(54) Title: QUINOLINE INHIBITORS OF cGMP PHOSPHODIESTERASE

(57) Abstract: Compounds of the formula (1) are useful as inhibitors of cGMP PDE especially Type 5.

QUINOLINE INHIBITORS OF cGMP PHOSPHODIESTERASE

Related Inventions

This application is related to, and pursuant to 35 U.S.C. § 119(e)
5 claims the benefit of priority of US application Serial No. 60/230,267, filed
September 6, 2000.

Field of the Invention

The present invention relates to quinoline compounds, to methods
10 of using such compounds in treating cGMP-associated conditions such as
erectile dysfunction, and to pharmaceutical compositions containing such
compounds.

Background of the Invention

15 Erectile dysfunction is the inability to obtain and maintain a penile
erection sufficient for sexual intercourse or other sexual expression. A
number of factors can place an individual at risk for this disorder, for example,
trauma, pelvic surgery, hypercholesterolemia, ischemic heart disease,
peripheral vascular disease, chronic renal failure, diabetes, the use of certain
20 medicaments including some types of antihypertensive agents, digoxin, or the
excessive use of narcotics, alcohol, tobacco, etc. Methods for treating
erectile dysfunction include the use of vacuum devices and penile implants,
as well as the administration of medicaments such as yohimbine, papaverine
and apomorphine. Improved methods for treating this disorder are sought,
25 however, as the aforementioned methods do not provide sufficient efficacy
and/or are accompanied by drawbacks or side effects such as erosion, pain,
priapism, or gastrointestinal discomfort.

A penile erection is dependent upon the presence of adequate levels
of cyclic guanosine 3',5'-monophosphate (cGMP), especially in corpora
30 cavernosa tissue. Thus, administering an inhibitor of a cGMP
phosphodiesterase (cGMP PDE), particularly a selective inhibitor of cGMP


PDE Type 5 (PDE 5), provides a means for achieving and maintaining an erection and therefore, for treating erectile dysfunction. See Trigo-Rocha *et al.*, "Nitric Oxide and cGMP: Mediators of Pelvic Nerve-Stimulated Erection in Dogs," *Am. J. Physiol.*, Vol. 264 (Feb. 1993); Bowman *et al.*, "Cyclic GMP Mediates Neurogenic Relaxation in the Bovine Retractor Penis Muscle," *Br. J. Pharmac.*, 81, 665-674 (1984); and Rajfer *et al.*, "Nitric Oxide as a Mediator of Relaxation of the Corpus Cavernosum in Response to Nonadrenergic, Noncholinergic Neurotransmission," *New England J. Med.*, 326, 2, 90-94 (Jan. 1992). Sildenafil, for example, has been described as a PDE 5 inhibitor useful for treating erectile dysfunction. See *Drugs of the Future*, 22, 138-143 (1997).

Recent examples of other compounds claimed as PDE 5 inhibitors include fused pyridazine compounds (WO 96/05176 and US Patent application Serial No. 09/393,833), anthranilic acid derivatives (US 5,716,993), fused pyridopyridazine compounds (US Patent application Serial No. 09/526,162), and quinazolinone compounds (US Patent No. 6,087,368).

The present invention provides compounds that are potent and selective inhibitors of cGMP PDE 5. These compounds may be employed in treating erectile dysfunction. In view of their activity, these compounds can also be used in treating other disorders responding to the inhibition of cGMP PDE, such as various cardiovascular disorders.

Summary of the Invention

The present invention provides quinoline compounds of the following formula (I) or salts thereof, for use as inhibitors of cGMP PDE, especially Type 5:



wherein:

- R_2 , R_3 , R_7 and R_8 are independently hydrogen, halogen, alkyl, substituted alkyl, alkoxy, nitro, cyano, aryl, heteroaryl, or heterocyclo;
- 5 R_3 is $-(CH_2)_zY$, wherein z is 0, 1, 2, or 3;
- R_4 and R_5 (i) are independently hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, or heteroaryl, with the proviso that R_4 and R_5 are not both hydrogen; or (ii) taken together form a heterocyclo ring;
- Y is selected (i) independently from $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$,
- 10 $-O(C=O)NR_{10}R_{11}$, $-NR_{10}R_{11}$, $-NR_{10}(C=O)NR_{11}R_{12}$, $-CH[(C=O)NR_{10}R_{11}]_2$, $-(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, $-S(O)_mR_9$, $-SO_2NR_{10}R_{11}$, imidazole, substituted imidazole, triazole, substituted triazole, or cyano, or (ii) together with one of R_4 and R_5 to form a heterocyclo ring therewith;
- m is 0, 1, or 2;
- 15 R_9 is hydrogen, alkyl, substituted alkyl, hydroxy, alkoxy, cycloalkyl, substituted cycloalkyl, heterocyclo, aryl, heteroaryl, or pentafluorophenyl; and
- R_{10} , R_{11} , and R_{12} (i) are independently selected from hydrogen, alkyl, substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl, heterocyclo, and heteroaryl; or (ii) taken together wherein R_{10} forms a
- 20 three- to seven-membered heterocyclo ring with R_{11} or R_{12} , or R_{11} forms a three- to seven-membered heterocyclo ring with R_{12} .

- The invention further provides pharmaceutical compositions adapted for use in treating cGMP-associated conditions comprising a pharmaceutically
- 25 acceptable diluent or carrier and at least one compound of the formula (I) or salt thereof, wherein R_2 and R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are as defined above and R_3 is selected from hydrogen and $-(CH_2)_zY$, with the proviso that at least one of R_2 , R_3 , R_6 , R_7 , and R_8 is not hydrogen. The invention further provides methods for treating cGMP-associated conditions
- 30 comprising administering to a mammal in need of such treatment a therapeutically-effective amount of one or more compounds of the formula (I)

or salt thereof, wherein R_2 and R_4 , R_5 , R_6 , R_7 , R_8 , R_9 , R_{10} , R_{11} and R_{12} are as defined above and R_3 is selected from hydrogen and $-(CH_2)_xY$, with the proviso that at least one of R_2 , R_3 , R_6 , R_7 , and R_8 is not hydrogen.

Detailed Description of the Invention

The following are definitions of terms used in this specification. The initial definition provided for a group or term herein applies to that group or term throughout the present specification, individually or as part of another group, unless otherwise indicated.

The term "alkyl" refers to straight or branched chain hydrocarbon groups having 1 to 12 carbon atoms, preferably 1 to 8 carbon atoms. Lower alkyl groups, that is, alkyl groups of 1 to 4 carbon atoms, are most preferred.

The term "substituted alkyl" refers to an alkyl group as defined above having one, two or three substituents selected from the group consisting of halo, amino, cyano, hydroxy, alkoxy, alkylthio, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -C(=O)H, -CO₂H, -CO₂-alkyl, cycloalkyl, substituted cycloalkyl, aryl, heteroaryl, or heterocycle. The term "substituted alkyl" also includes an alkyl group as defined above substituted with N(substituted alkyl) or N(substituted alkyl)₂, or in other words, the groups (CH₂)_nNHR' and (CH₂)_nNR'R", wherein each of R' and R" comprises a substituted alkyl or R' and R" together form a heterocyclo ring.

The term "alkoxy" refers to an alkyl group as defined above bonded through an oxygen (-O-). The term "alkylthio" refers to an alkyl group as defined above bonded through a sulfur (-S-).

The term "cycloalkyl" refers to fully saturated and partially unsaturated hydrocarbon rings of 3 to 9, preferably 3 to 7, carbon atoms as well as such rings having a fused aryl ring such as indan.

The term "substituted cycloalkyl" refers to such rings having one, two or three substituents, preferably one, selected from the group consisting of alkyl, substituted alkyl, alkoxy, alkylthio, halo, hydroxy, cyano, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -CO₂H, -CO₂-lower alkyl, aryl, heterocyclo, heteroaryl, keto, =N-OH, =N-O-lower alkyl, and a five or six membered ketal, *i.e.* 1,3-dioxolane or 1,3-dioxane.

The term "halo" refers to chloro, bromo, fluoro and iodo.

- The term "aryl" refers to phenyl, 1-naphthyl and 2-naphthyl, with phenyl being preferred. The term "aryl" includes such rings having from zero, one, two or three substituents, selected from the group consisting of alkyl, substituted alkyl, alkoxy, alkylthio, halo, hydroxy, nitro, cyano, amino, -
- 5 NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -CO₂H, -(C=O)alkyl, -CO₂-alkyl, cycloalkyl, substituted cycloalkyl, -(C=O)NH₂, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -(C=O)N(alkyl)₂, -NH-CH₂-CO₂H, -NH-CH₂-CO₂-alkyl, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, heterocyclo, and heteroaryl.
- 10 The term "heterocyclo" refers to substituted and unsubstituted non-aromatic 3 to 7 membered monocyclic groups, 7 to 11 membered bicyclic groups, and 10 to 15 membered tricyclic groups which have at least one heteroatom (O, S or N) in at least one of the rings. Each ring of the heterocyclo group containing a heteroatom can contain one or two oxygen or
- 15 sulfur atoms and/or from one to four nitrogen atoms provided that the total number of heteroatoms in each ring is four or less, and further provided that the ring contains at least one carbon atom. The fused rings completing the bicyclic and tricyclic groups may contain only carbon atoms and may be saturated, partially saturated, or unsaturated. The nitrogen and sulfur atoms
- 20 may optionally be oxidized and the nitrogen atoms may optionally be quaternized. The heterocyclo group may be attached at any available nitrogen or carbon atom. The heterocyclo ring may contain one, two or three substituents selected from the group consisting of halo, amino, cyano, alkyl, substituted alkyl, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, alkoxy, alkylthio,
- 25 hydroxy, nitro, phenyl, benzyl, phenylethyl, phenyloxy, phenylthio, -CO₂H, -CO₂-alkyl, cycloalkyl, substituted cycloalkyl, -(C=O)NH₂, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -(C=O)N(alkyl)₂, -NH-CH₂-CO₂H, -NH-CH₂-CO₂-alkyl, heterocyclo, heteroaryl, keto, =N-OH, =N-O-lower alkyl, and a five or six membered ketal, *i.e.*, 1,3-dioxolane or 1,3-dioxane.
- 30 Exemplary monocyclic groups include azetidiny, pyrrolidiny, oxetany, imidazoliny, oxazolidiny, isoxazolidiny, thiazolidiny, isothiazolidiny,

tetrahydrofuranyl, piperidinyl, piperazinyl, 2-oxopiperazinyl, 2-oxopiperidinyl, 2-oxopyrrolodinyl, 2-oxoazepinyl, azepinyl, 4-piperidinyl, tetrahydropyranyl, morpholinyl, thiamorpholinyl, thiamorpholinyl sulfoxide, thiamorpholinyl sulfone, 1,3-dioxolane and tetrahydro-1,1-dioxothieryl and the like.

- 5 Exemplary bicyclic heterocyclo groups include quinuclidinyl.

The term "heteroaryl" refers to substituted and unsubstituted aromatic
5 or 6 membered monocyclic groups, 9 or 10 membered bicyclic groups, and
11 to 14 membered tricyclic groups which have at least one heteroatom (O, S
or N) in at least one of the rings. Each ring of the heteroaryl group containing
10 a heteroatom can contain one or two oxygen or sulfur atoms and/or from one
to four nitrogen atoms provided that the total number of heteroatoms in each
ring is four or less and each ring has at least one carbon atom. The fused
rings completing the bicyclic and tricyclic groups may contain only carbon
atoms and may be saturated, partially saturated, or unsaturated. The
15 nitrogen and sulfur atoms may optionally be oxidized and the nitrogen atoms
may optionally be quaternized. Heteroaryl groups which are bicyclic or
tricyclic must include at least one fully aromatic ring but the other fused ring
or rings may be aromatic or non-aromatic. The heteroaryl group may be
attached at any available nitrogen or carbon atom of any ring. The heteroaryl
20 ring system may contain one, two or three substituents selected from the
group consisting of halo, amino, cyano, alkyl, substituted alkyl, -NH(alkyl), -
NH(cycloalkyl), -N(alkyl)₂, alkoxy, alkylthio, hydroxy, nitro, phenyl, benzyl,
phenylethyl, phenoxy, phenylthio, -CO₂H, -CO₂-alkyl, cycloalkyl, substituted
cycloalkyl, -(C=O)NH₂, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -
25 (C=O)N(alkyl)₂, -NH-CH₂-CO₂H, -NH-CH₂-CO₂-alkyl, heterocylco, and
heteroaryl.

Exemplary monocyclic heteroaryl groups include pyrrolyl, pyrazolyl,
pyrazolinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, isothiazolyl,
furanyl, thienyl, oxadiazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl
30 and the like.

Exemplary bicyclic heteroaryl groups include indolyl, benzothiazolyl, benzodioxolyl, benzoxazolyl, benzothienyl, quinolinyl, tetrahydroisoquinolinyl, isoquinolinyl, benzimidazolyl, benzopyranyl, indolizynyl, benzofuranyl, chromonyl, coumarinyl, benzopyranyl, cinnolynyl, quinoxalinyl, indazolyl, 5 pyrolopyridyl, furopyridinyl, dihydroisoindolyl, tetrahydroquinolinyl and the like.

Exemplary tricyclic heteroaryl groups include carbazolyl, benzidolyl, phenanthrolyl, acridinyl, phenanthridinyl, xanthenyl and the like.

The term "substituted imidazole" refers to an imidazole, an aryl-fused 10 imidazole such as benzimidazole, or a heteroaryl-fused imidazole such as a pyridoimidazole which contain one or two substituents selected from the group consisting of hydrogen, alkyl, substituted alkyl, alkoxy, alkylthio, halo, hydroxy, nitro, cyano, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -CO₂H, -CO₂-alkyl, cycloalkyl, substituted cycloalkyl, -(C=O)NH₂, -(C=O)NH(alkyl), - 15 (C=O)NH(cycloalkyl), -(C=O)N(alkyl)₂, -NH-CH₂-CO₂H, -NH-CH₂-CO₂-alkyl, phenyl, benzyl, phenylethyl, phenoxy, phenylthio, heterocyclo, and heteroaryl.

The term "substituted triazole" refers to a triazole having at least one substituent selected from the group consisting of alkyl, substituted alkyl, 20 alkoxy, alkylthio, halo, hydroxy, nitro, cyano, amino, -NH(alkyl), -NH(cycloalkyl), -N(alkyl)₂, -CO₂H, -CO₂-alkyl, cycloalkyl, substituted cycloalkyl, -(C=O)NH₂, -(C=O)NH(alkyl), -(C=O)NH(cycloalkyl), -(C=O)N(alkyl)₂, -NH-CH₂-CO₂H, -NH-CH₂-CO₂-alkyl, phenyl, benzyl, phenylethyl, phenoxy, phenylthio, heterocyclo, and heteroaryl.

25 Throughout the specification, groups and substituents thereof may be chosen to provide stable moieties and compounds.

The compounds of formula I form salts which are also within the scope of this invention. Reference to a compound of the formula I herein is understood to include reference to salts thereof, unless otherwise indicated.

30 The term "salt(s)", as employed herein, denotes acidic and/or basic salts formed with inorganic and/or organic acids and bases. In addition, when a

compound of formula I contains both a basic moiety, such as, but not limited to an amine or a pyridine or imidazole ring, and an acidic moiety, such as, but not limited to a carboxylic acid, zwitterions ("inner salts") may be formed and are included within the term "salt(s)" as used herein. Pharmaceutically acceptable (*i.e.*, non-toxic, physiologically acceptable) salts are preferred, although other salts are also useful, *e.g.*, in isolation or purification steps which may be employed during preparation. Salts of the compounds of the formula I may be formed, for example, by reacting a compound of the formula I with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.

The compounds of formula I which contain a basic moiety, such as, but not limited to an amine or a pyridine or imidazole ring, may form salts with a variety of organic and inorganic acids. Exemplary acid addition salts include acetates (such as those formed with acetic acid or trihaloacetic acid, for example, trifluoroacetic acid), adipates, alginates, ascorbates, aspartates, benzoates, benzenesulfonates, bisulfates, borates, butyrates, citrates, camphorates, camphorsulfonates, cyclopentanepropionates, digluconates, dodecylsulfates, ethanesulfonates, fumarates, glucoheptanoates, glycerophosphates, hemisulfates, heptanoates, hexanoates, hydrochlorides (formed with hydrochloric acid), hydrobromides (formed with hydrogen bromide), hydroiodides, 2-hydroxyethanesulfonates, lactates, maleates (formed with maleic acid), methanesulfonates (formed with methanesulfonic acid), 2-naphthalenesulfonates, nicotines, nitrates, oxalates, pectinates, persulfates, 3-phenylpropionates, phosphates, picrates, pivalates, propionates, salicylates, succinates, sulfates (such as those formed with sulfuric acid), sulfonates (such as those mentioned herein), tartrates, thiocyanates, toluenesulfonates such as tosylates, undecanoates, and the like.

The compounds of formula I which contain an acidic moiety, such as, but not limited to a carboxylic acid, may form salts with a variety of organic

and inorganic bases. Exemplary basic salts include ammonium salts, alkali metal salts such as sodium, lithium, and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salts with organic bases (for example, organic amines) such as benzathines, dicyclohexylamines, 5 hydrabamines [formed with N,N-bis(dehydro-abietyl)ethylenediamine], N-methyl-D-glucamines, N-methyl-D-glucamides, t-butyl amines, and salts with amino acids such as arginine, lysine and the like. Basic nitrogen-containing groups may be quaternized with agents such as lower alkyl halides (e.g., methyl, ethyl, propyl, and butyl chlorides, bromides and iodides), dialkyl 10 sulfates (e.g., dimethyl, diethyl, dibutyl, and diamyl sulfates), long chain halides (e.g., decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides), aralkyl halides (e.g., benzyl and phenethyl bromides), and others.

Prodrugs and solvates of the compounds of this invention are also contemplated herein. The term "prodrug", as employed herein, denotes a 15 compound which, upon administration to a subject, undergoes chemical conversion by metabolic or chemical processes to yield a compound of the formula I, and/or a salt and/or solvate thereof. Solvates of the compounds of formula I are preferably hydrates.

Compounds of the formula I, and salts thereof, may exist in their 20 tautomeric form (for example, as an amide or imino ether). All such tautomeric forms are contemplated herein as part of the present invention.

All stereoisomers of the present compounds, such as those, for example, which may exist due to asymmetric carbons on the R_2 to R_{12} substituents, including enantiomeric forms (which may exist even in the 25 absence of asymmetric carbons) and diastereomeric forms, are contemplated and within the scope of this invention. Individual stereoisomers of the compounds of this invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other or other selected, stereoisomers. The chiral centers of the present invention 30 can have the S or R configuration as defined by the IUPAC 1974 Recommendations.


Methods of Preparation

The compounds of the present invention may be prepared by methods such as those illustrated in the following Schemes I to X. Starting materials
5 are commercially available or can be readily prepared by one of ordinary skill in the art using known methods. For all of the schemes and compounds, the groups R_2 , R_3 , R_4 , R_5 , R_6 , R_7 , and R_8 are as described above for a compound of formula I, unless otherwise indicated.

Solvents, temperatures, pressures, and other reaction conditions may
10 readily be selected by one of ordinary skill in the art. For example, in these schemes exemplary hydroxide sources may include sodium hydroxide or lithium hydroxide; an exemplary reducing reagent and inert solvent (for reducing a carboxylic acid or ester group to an alcohol) includes lithium tri-*t*-butoxyaluminumhydride and tetrahydrofuran (THF); exemplary dehydrating/
15 chlorinating agents include POCl_3 , PCl_5 , SOCl_2 or oxalyl chloride; exemplary leaving groups (LG) include triflate, mesylate, tosylate, or halide; and exemplary reagents (for converting a hydroxyl group to a leaving group) include trifluoromethanesulfonyl chloride, toluenesulfonyl chloride, methanesulfonyl chloride, phosphorus oxychloride, thionyl chloride, and
20 phosphorus pentachloride. Exemplary solvents, as appropriate, may be selected from 1,2-dichlorobenzene, methylene chloride, dimethylformamide (DMF), alcohols, ethers, including diphenyl ether, tetrahydrofuran and dioxane, N,N-dimethylformamide, and acetonitrile, water, mixtures of ethers and water, and the like.

25 High Speed Analoging (HSA) may be employed in the preparation of compounds.

SCHEME I



5

Compounds of formula Ia wherein R₃ is hydrogen can be prepared via the decarboxylation of a compound of formula Ib in an appropriate degassed inert solvent (e.g., 1,2-dichlorobenzene and diphenyl ether) at elevated temperature.

10

Compounds of formula Ib wherein R₃ is -CO₂H can be prepared by the hydrolysis of compounds of formula Ic using a hydroxide source and appropriate solvent (e.g., water, alcohols, and mixtures of ethers and water).

15

Compounds of formula Id wherein R₃ is -CH₂OH can be prepared by reducing a compound of formula Ib or Ic with an appropriate reducing reagent in an inert (reaction) solvent.

Compounds of formula Ic wherein R₃ is -CO₂R₉ can be prepared by reacting compounds of formula II with an amine of the formula NHR₄R₅. The reaction may be performed in a solvent as appropriate, such as an alcohol, in

the presence of an appropriate base, such as triethylamine, and typically under elevated temperatures.

Compounds of formula **II** can be prepared by reacting compounds of formula **III** with an appropriate dehydrating/chlorinating agent, typically under elevated temperatures.


Compounds of formula **III** can be prepared from compounds of formula **IV** via an intramolecular cyclization typically under elevated temperatures in an inert solvent, as appropriate, or in neat form.

Compounds of formula **IV** can be prepared by combining compounds of formula **V** and **VI** either neat or in an inert solvent as appropriate, typically under elevated temperatures.

Compounds of formula **V** and formula **VI** are either commercially available or available via methods known to one skilled in the art.

15

SCHEME II



20

Compounds of formula **Ia** wherein R_2 is halogen and R_3 is $-\text{CH}_2\text{OH}$ can be prepared by reducing a compound of formula **Ib** or **Ic** with an appropriate reducing reagent in an inert (reaction) solvent.

Compounds of formula **1f** wherein R₂ is halogen and R₃ is -CO₂H can be prepared by the hydrolysis of compounds of formula **1g** using a hydroxide source in appropriate solvent.


Compounds of formula **Ig** wherein R_2 is halogen and R_3 is $-\text{CO}_2R_9$ (R_9 is
5 other than hydrogen) can be prepared by reacting compounds of formula **VII**
with an amine of the formula NHR_6 (as in Scheme I for formula **Ic**).

Compounds of formula **VII** can be prepared by reacting compounds of formula **VIII** with an appropriate dehydrating /chlorinating agent typically under elevated temperatures.

10 Compounds of formula **VIII** can be prepared from compounds of formula **IX** by a condensation with a malonate derivative using base in an appropriate solvent. Sodium alkoxides are exemplary bases and alcohols exemplary solvents.

Compounds of formula IX are either commercially available or
15 available via methods known to one skilled in the art.

SCHEME III



Compounds of formula **Ih** wherein R_3 is $-(CH_2)_2OH$ can be prepared by reducing a compound of formula **Ii** with an appropriate reducing reagent in an inert solvent.

- 5 Compounds of formula **Ii** wherein R_3 is $-CH_2CO_2H$ can be prepared by the hydrolysis of compounds of formula **Ij** using a hydroxide source in appropriate solvent.

- 10 Compounds of formula **Ij** wherein R_3 is $-CH_2CN$ can be prepared via the displacement of the leaving group (LG) from a compound of formula **X** using an appropriate nucleophile in an inert solvent (*e.g.*, methylene chloride). Nucleophiles may include cyanides from HCN, KCN, or NaCN, or tetrabutylammonium cyanide.

- 15 Compounds of formula **X** can be prepared via reaction of compounds of formula **Id** or **Ie** with an appropriate reagent which converts the hydroxyl group to a leaving group (LG) in an inert solvent (*e.g.*, methylene chloride).


Compounds of formula **Id** or **Ie** wherein R_3 is $-CH_2OH$, can be prepared by the methods described above in Schemes I and II.

- 20 Compounds of formula **Ik** wherein R_3 is $-(CH_2)_3OH$ can be prepared by reducing a compound of formula **II** with an appropriate reducing reagent in an inert solvent. Compounds of formula **II** can be prepared by decarboxylation of compounds of formula **XI** at elevated temperature in an inert solvent (*e.g.*, DMF).

- 25 Compounds of formula **XI** can be prepared by the hydrolysis of compounds of formula **Im** using a hydroxide source and an appropriate solvent.

Compounds of formula **Im** wherein R_3 is $-CH_2CH(CO_2R_9)_2$ can be prepared via the displacement of the leaving group (LG) from a compound of formula **X** using an appropriate nucleophile (*e.g.*, alkyl malonate) in an inert solvent (*e.g.*, methylene chloride).

SCHEME IV




5

Compounds of formula **In** wherein R_3 is $-(CH_2)_zY$ can be prepared via the displacement of the leaving group (LG) from a compound of formula **XI** using an appropriate nucleophile in an inert solvent. Exemplary nucleophiles include alcohols of the formula HOR_9 , amines of the formula $HNR_{10}R_{11}$, mercaptans of the formula HSR_9 , cyanides as in Scheme III, imidazole, substituted imidazoles, triazole, and substituted triazoles.

Compounds of formula **XII** can be prepared via reaction of compounds of formulas **Id**, **le**, **lh**, **lk** with an appropriate reagent which converts the hydroxyl group to a leaving group (LG) in an appropriate inert solvent.

Compounds of formulas **Id** ($z=1$), **le** ($z=1$), **lh** ($z=2$), **lk** ($z=3$) can be prepared using the methods described above in Schemes I, II or III.

SCHEME V



20


Compounds of formula **1o** wherein R_3 is $-(CH_2)_z(C=O)NR_{10}R_{11}$ can be prepared via the aminolysis of an active ester (AE) of formula **XIII** using an amine of the formula $NHR_{10}R_{11}$ in an inert solvent (e.g., ethers, described above, and methylene chloride).

Compounds of formula **1p** wherein R_3 is $-(CH_2)_zCO_2R_9$ can be prepared via the esterification of a compound of formula **XIII** using an alcohol of the formula HOR_9 , with the alcohol used as solvent or in an inert solvent such as ethers, described above, or methylene chloride.

Compounds of formula **XIII** can be prepared via the activation of the carboxylic acid in compounds of formula **1q** using an appropriate carboxylic acid activating agent in an appropriate solvent. Exemplary activating agents include carbonyldiimidazole or dicyclohexyl carbodiimide and pentafluorophenol.

Compounds of formula **1q** wherein R_3 is $-(CH_2)_zCO_2H$ can be prepared with the methods described in Schemes III and IV.

SCHEME VI



Compounds of formula **1r** wherein R_3 is $-(CH_2)_zSO_2NR_{10}R_{11}$ can be prepared via the aminolysis of a sulfonyl chloride of formula **XIV** using an


amine of the formula $\text{NHR}_{10}\text{R}_{11}$ in an inert solvent (e.g., ethers or methylene chloride).

Compounds of formula **XIV** can be prepared by reacting compounds of formula **Is** with an appropriate dehydrating/chlorinating agent typically under elevated temperatures.

Compounds of formula **Is** wherein R_3 is $-(\text{CH}_2)_z\text{SO}_2\text{OH}$ can be prepared by oxidation of compounds of formula **It**.

Compounds of formula **It** wherein R_3 is $-(\text{CH}_2)_z\text{SH}$ can be prepared by the methods described above in Scheme IV.

SCHEME VII




Compounds of formula **Iu** wherein R_3 is $-(\text{CH}_2)_z\text{NR}_{10}(\text{C}=\text{O})\text{OR}_{11}$ can be prepared by reacting compounds of formula **Iv** with a chloroformate in an inert solvent.

Compounds of formula **Iw** wherein R_3 is $-(\text{CH}_2)_z\text{NR}_{10}(\text{C}=\text{O})\text{NR}_{11}\text{R}_{12}$ can be prepared by reacting compounds of formula **Ix** with an alkylating agent under basic conditions in an inert solvent.

Compounds of formula **Ix** wherein R_3 is $-(CH_2)_zNR_{10}(C=O)NHR_{11}$ can be prepared by reacting compounds of formula **Iv** with an isocyanate in an inert solvent.

Compounds of formula **Iv** can be prepared with the methods described
5 above in Scheme IV.

SCHEME VIII



10


Compounds of formula **Iy** wherein R_3 and R_5 form a heterocyclo carboxamide ring can be prepared from compounds of formula **Iz** by hydrolysis and decarboxylation.

Compounds of formula **Iz** wherein R_3 and R_5 form a 2-carboxylate heterocyclo carboxamide ring can be prepared from compounds of formula **Iaa** by a base effected cyclization.

Compounds of formula **Iaa** wherein R_3 is $-(CH_2CH(CO_2R_9))_2$ and R_5 is hydrogen can be prepared with the methods described in Scheme III.

20

SCHEME IX



5


Compounds of formula **lab** wherein R_3 is $-(CH_2)_z$ OR $OR_9(OR_{10})NR_{11}R_{12}$ can be prepared from compounds of formula **lac** by a base-effected alkylation.

Compounds of formula **lac** wherein R_3 is $-(CH_2)_z$ OR $OR_9(OH)NR_{11}R_{12}$ can be prepared from compounds of formula **lab** by an aminolysis of the epoxide.

10

Compounds of formula **lad** wherein R_3 is $-(CH_2)_z$ OR $OR_9NR_{10}R_{11}$ can be prepared by the method described in Scheme IV.

SCHEME X



15

Compounds of formula **Ia** wherein R_3 is hydrogen can also be prepared from compounds of formula **XV** by a base-effected alkylation.

Compounds of formula **XV** can be prepared from compounds of formula **XVI** by a Curtius rearrangement.

- 5 Compounds of formula **XVI** can be prepared by reacting compounds of formula **XVII** with an appropriate methyl ketone.

Compounds of formula **XVII** are either commercially available or available via methods known to one skilled in the art.

10

Preferred Compounds

Preferred compounds of this invention are those of formula (I) and/or pharmaceutically acceptable salts thereof having the following definitions:



15

wherein,

R_2 , R_8 , and R_7 are independently hydrogen, halogen, alkyl, substituted alkyl, nitro, cyano, aryl, or heteroaryl;

R_9 is hydrogen, alkyl, or substituted alkyl, including $-(CH_2)_nNR_{13}R_{14}$, wherein

- 20 R_{13} and R_{14} (i) are independently selected from hydrogen, alkyl, or substituted alkyl, or (ii) taken together form a heterocyclo ring;

R_3 is $-(CH_2)_zY$;

R_4 is hydrogen, lower alkyl, or forms a heterocyclo ring with Y or R_3 ;

R_5 is substituted alkyl;

- 25 R_6 is hydrogen, halogen, trifluoromethyl, nitro, cyano, or aryl;

Y is (i) independently selected from $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$,

$-O(C=O)NR_{10}R_{11}$, $-NR_{10}R_{11}$, $-NR_{10}(C=O)NR_{11}R_{12}$, $-CH[(C=O)NR_{10}R_{11}]_2$, -

$(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, $-S(O)_mR_9$, $-SO_2NR_{10}R_{11}$, imidazole,

- substituted imidazole, triazole, substituted triazole, or cyano, or (ii) taken together with R_4 or R_5 , forming a heterocyclic ring therewith;
- m is 0, 1, or 2;
- n is 0, 1, 2, or 3;
- 5 z is 0, 1, 2, or 3;
- R_9 is hydrogen, alkyl, substituted alkyl, hydroxy, alkoxy, cycloalkyl, substituted cycloalkyl, heterocyclo, aryl, heteroaryl, or pentafluorophenyl; and
- R_{10} , R_{11} , and R_{12} (i) are independently selected from hydrogen, alkyl, substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl,
- 10 heterocyclo, or heteroaryl; or (ii) taken together wherein R_{10} with R_{11} or R_{12} forms a three- to seven-membered heterocyclic ring, or R_{10} with R_{12} forms a three- to seven-membered heterocyclic ring.

- More preferred are the compounds of formula (I), above, and/or
- 15 pharmaceutically acceptable salts thereof, wherein
- R_2 is hydrogen, halogen, lower alkyl, or pyridine;
- R_3 is $-(CH_2)_2Y$;
- R_4 is hydrogen, methyl, or forms a heterocyclic ring with Y or R_3 ;
- R_5 is substituted alkyl, wherein said substituted alkyl comprises an aryl,
- 20 cycloalkyl, or heteroaryl substituent;
- R_8 is hydrogen, halogen, trifluoromethyl, or cyano;
- R_7 is hydrogen or trifluoromethyl;
- R_8 is hydrogen, alkyl, substituted alkyl, or $-(CH_2)_nNR_{13}R_{14}$, wherein R_{13} and R_{14} (i) are independently selected from hydrogen, alkyl, or substituted
- 25 alkyl, or (ii) together form a heterocyclic ring;
- Y is (i) $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$, $-OR_9NR_{10}R_{11}$, $-NR_{10}R_{11}$, $-(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, or (ii) together with R_4 forms a heterocyclic ring;
- n is 0, 1, 2, or 3;
- 30 z is 0, 1, 2, or 3;

- R_9 is hydrogen, alkyl, substituted alkyl, alkoxy, heterocyclo, or pentafluorophenyl;
- R_{10} and R_{11} are (i) independently selected from hydrogen, alkyl, substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl, heterocyclo, or heteroaryl; or (ii) taken together wherein R_{10} and R_{11} forms a three- to seven-membered heterocyclo ring; and
- R_{12} is aryl, cycloalkyl, or heteroaryl.

- Advantageously, R_5 comprises cycloalkyl or an alkyl substituted with aryl. When R_5 is an alkyl substituted with an aryl, advantageously the aryl has one to two substituents wherein at least one of the substituents is selected from halogen (e.g., chloro, bromo, fluoro), alkoxy (e.g., methoxy), or a lower alkyl. Advantageously, when R_9 comprises a heterocyclo ring with a nitrogen heteroatom, said nitrogen heteroatom has a substituent X_1 selected from lower alkyl, substituted alkyl, and cycloalkyl. When R_4 and R_3 (or Y) form a heterocyclo ring, said ring advantageously is unsubstituted or has at least one substituent X_2 comprising CO_2 (alkyl).

- Most preferred are the compounds of formula (I) and/or pharmaceutically acceptable salts thereof, wherein:

- R_2 is hydrogen or chloro;
- R_3 is $-(CH_2)_zY$, wherein z is 0, 1, 2, or 3;
- R_4 is hydrogen;
- R_5 is 3-chloro-4-methoxyphenylmethyl;
- R_6 is cyano;
- R_7 is hydrogen;
- R_8 is hydrogen, alkyl, or substituted alkyl;
- Y is $-OR_9$, $-NR_{10}R_{11}$, $-CO_2R_9$, or $-(C=O)NR_{10}R_{11}$;
- R_9 is hydrogen, alkyl, or substituted alkyl; and


R_{10} and R_{11} (i) are each independently hydrogen, alkyl, substituted alkyl, aryl, heterocyclo, or heteroaryl; or (ii) together form a five- to seven-membered heterocyclo ring.

5

**Preferred Pharmaceutical Compositions for
Treating cGMP-Associated Conditions**

Preferred pharmaceutical compositions of this invention are those compositions adapted for use in treating cGMP-associated conditions comprising a pharmaceutically acceptable diluent or carrier and at least one

10 compound of the formula (I) and/or pharmaceutically acceptable salts thereof:



wherein,

- 15 R_2 , R_9 , and R_7 are independently hydrogen, halogen, alkyl, substituted alkyl, nitro, cyano, aryl, or heteroaryl;
 R_8 is hydrogen, alkyl, substituted alkyl, or $-(CH_2)_nNR_{13}R_{14}$, wherein R_{13} and R_{14} (i) are independently selected from hydrogen, alkyl, or substituted alkyl, or (ii) taken together form a heterocyclo ring;
- 20 R_3 is hydrogen or $-(CH_2)_2Y$;
 R_4 is hydrogen, lower alkyl, or forms a heterocyclo ring with Y or R_3 ;
 R_5 is substituted alkyl;
 R_6 is hydrogen, halogen, trifluoromethyl, nitro, cyano or aryl;
Y is selected (i) independently from $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$, $-O(C=O)NR_{10}R_{11}$, $-NR_{10}R_{11}$, $-NR_{10}(C=O)NR_{11}R_{12}$, $-CH[(C=O)NR_{10}R_{11}]_2$, $-(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, $-S(O)_mR_9$, $-SO_2NR_{10}R_{11}$, imidazole, substituted imidazole, triazole, substituted triazole, or cyano, or (ii) together with R_4 or R_5 , forming a heterocyclo ring therewith;
- 25

m is 0, 1, or 2;

n is 0, 1, 2, or 3;

z is 0, 1, 2, or 3;

- R_9 is hydrogen, alkyl, substituted alkyl, hydroxy, alkoxy, cycloalkyl, substituted
 5 cycloalkyl, heterocyclo, aryl, heteroaryl, or pentafluorophenyl; and
 R_{10} , R_{11} , and R_{12} (i) are independently selected from hydrogen, alkyl,
 substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl,
 heterocyclo, or heteroaryl; or (ii) taken together wherein R_{10} with R_{11} or
 R_{12} forms a three- to seven-membered heterocyclo ring, or R_{10} with R_{12}
 10 forms a three- to seven-membered heterocyclo ring.

- In the pharmaceutical compositions wherein R_3 of formula (I) is
 hydrogen, it is preferred that R_3 is cyano. In an alternative embodiment
 where R_3 is hydrogen, R_2 advantageously is selected from heteroaryl
 15 including pyridine. Advantageously, when R_3 is hydrogen, R_2 does not include
 hydrogen, chlorine, or methyl.

- More preferred pharmaceutical compositions are those including a
 pharmaceutically acceptable diluent or carrier and at least one compound of
 the formula (I) and/or pharmaceutically acceptable salts thereof, wherein
 20 R_2 is hydrogen, halogen, lower alkyl, or pyridine;
 R_3 is hydrogen or $-(CH_2)_zY$;
 R_4 is hydrogen, methyl, or forms a heterocyclo ring with Y or R_3 ;
 R_5 is alkyl substituted with aryl, cycloalkyl, or heterocyclo;
 R_6 is hydrogen, halogen, trifluoromethyl, or cyano;
 25 R_7 is hydrogen or trifluoromethyl;
 R_8 is hydrogen, alkyl, substituted alkyl, or $-(CH_2)_nNR_{13}, R_{14}$, wherein R_{13} and
 R_{14} (i) are independently selected from hydrogen, alkyl, or substituted
 alkyl, (ii) together form a heterocyclo ring;
 Y is (i) $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$, $-OR_9NR_{10}R_{11}$, $-NR_{10}R_{11}$,
 30 $-(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, or (ii) together with R_4 forms a
 heterocyclo ring;

n is 0, 1, 2, or 3;

z is 0, 1, 2, or 3;

R₉ is hydrogen, alkyl, substituted alkyl, alkoxy, heterocyclo, or pentafluorophenyl;

- 5 R₁₀ and R₁₁ are (i) independently selected from hydrogen, alkyl, substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl, heterocyclo, or heteroaryl; or (ii) taken together wherein R₁₀ and R₁₁ forms a three- to seven-membered heterocyclo ring; and

R₁₂ is aryl, cycloalkyl, or heteroaryl.

10

- Advantageously, R₆ comprises an alkyl substituted with cycloalkyl or an aryl group. When R₆ is alkyl substituted with aryl, advantageously the aryl has one to two substituents wherein at least one of the substituents is selected from halogen (*e.g.*, chloro, bromo, fluoro), alkoxy (*e.g.*, methoxy), or a lower
- 15 alkyl. Advantageously, when R₆ comprises a heterocyclo ring with a nitrogen heteroatom, said nitrogen heteroatom has a substituent X₁ selected from lower alkyl, substituted alkyl, and cycloalkyl. When R₄ and R₃ (or Y) form a heterocyclo ring, said ring advantageously is unsubstituted or has at least one substituent X₂ comprising CO₂(alkyl).

20

- Most preferred pharmaceutical compositions are those including a pharmaceutically acceptable diluent or carrier and at least one compound of the formula (I) and/or pharmaceutically acceptable salts thereof, wherein
- R₂ is hydrogen or chloro;

- 25 R₃ is -(CH₂)_zY, wherein z is 0, 1, 2, or 3;

R₄ is hydrogen;

R₅ is 3-chloro-4-methoxyphenylmethyl;

R₆ is cyano;

R₇ is hydrogen;

- 30 R₈ is hydrogen, alkyl or substituted alkyl;

Y is -OR₉, -NR₁₀R₁₁, -CO₂R₉, -(C=O)NR₁₀R₁₁;

R₉ is hydrogen, alkyl, or substituted alkyl; and
R₁₀ and R₁₁ (i) are each independently hydrogen, alkyl, substituted alkyl, aryl,
heterocyclo, or heteroaryl; or (ii) together form a five- to seven-
membered heterocyclo ring.

5

Utility

The compounds and compositions of this invention inhibit cGMP PDE,
and in particular are potent and selective inhibitors of cGMP PDE 5. Thus,
10 these compounds and compositions are useful in treating cGMP-associated
conditions. A "cGMP-associated condition", as used herein, denotes a
disorder which can be treated by inhibiting cGMP PDE or elevating the level
of cGMP in a subject, wherein treatment comprises prevention, partial
alleviation, or cure of the disorder. Inhibition of cGMP PDE or elevation of the
15 cGMP level may occur locally, for example, within certain tissues of the
subject, or more extensively throughout the subject being treated for such a
disorder. Treatment may be facilitated wherein elevation of the cGMP level
potentiates additional beneficial therapeutic effects, such as where elevation
of the cGMP level potentiates the effects of endothelium-derived relaxing
20 factor.

The inventive compounds and compositions are useful for treating a
variety of cardiovascular diseases including, but not limited to, hypertension,
angina (stable, unstable, and variant), (congestive) heart failure, restenosis,
atherosclerosis, and dyslipidemia, as well as reduced blood vessel patency,
25 thrombus, both venous and arterial, myocardial infarction, peripheral vascular
disease, stroke, bronchitis, chronic asthma, allergic asthma, allergic rhinitis,
glaucoma, benign prostate hyperplasia (BPH), and forms of cancer responsive
to the inhibition of cGMP PDE. In addition, these compounds are useful in
treating sexual dysfunction in both men (erectile dysfunction, for example,
30 due to diabetes mellitus, spinal cord injury, radical prostatectomy,
psychogenic etiology or any other cause) and women by improving blood flow

to the genitalia, especially, the corpus cavernosum. The compounds and compositions of this invention also are useful in treating diabetes mellitus and related conditions, and diseases of the gastrointestinal tract, such as those characterized by disorders of gut motility, including gastric paresis.

5 The present invention thus provides methods for treating cGMP-associated conditions, comprising administering to a subject in need thereof an effective amount of at least one compound of the formula I or a salt thereof, and/or pharmaceutical compositions as described above. Other therapeutic agents such as those described below may be employed in
10 combination with the compounds of formula I. In the methods of the present invention, such other therapeutic agent(s) may be administered prior to, simultaneously with, or following the administration of the inventive compound(s) and compositions.

 The present invention also provides pharmaceutical compositions
15 capable of treating a cGMP-associated condition, as described above. The compositions of the present invention may contain other therapeutic agents as described below, and may be formulated, for example, by employing conventional solid or liquid vehicles or diluents, as well as pharmaceutical additives of a type appropriate to the mode of desired administration (for
20 example, excipients, binders, preservatives, stabilizers, flavors, etc.) according to techniques such as those well known in the art of pharmaceutical formulation.

 The compounds and compositions of formula I may be administered by any suitable means, for example, orally, such as in the form of tablets,
25 capsules, granules or powders; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular or intrasternal injection or infusion techniques (e.g., as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally such as by inhalation spray; topically, such as in the form of a cream or ointment; rectally such as in the form of suppositories; or
30 liposomally; in dosage unit formulations containing non-toxic, pharmaceutically acceptable vehicles or diluents. These compounds may, for

example, be administered in a form suitable for immediate release or extended release. Immediate release or extended release may be achieved by the use of suitable pharmaceutical compositions or, particularly in the case of extended release, by the use of devices such as subcutaneous implants or osmotic pumps.

Exemplary compositions for oral administration include suspensions which may contain, for example, microcrystalline cellulose for imparting bulk, alginic acid or sodium alginate as a suspending agent, methylcellulose as a viscosity enhancer, and sweeteners or flavoring agents such as those known in the art; and immediate release tablets which may contain, for example, microcrystalline cellulose, dicalcium phosphate, starch, magnesium stearate and/or lactose and/or other excipients, binders, extenders, disintegrants, diluents and lubricants such as those known in the art. The compounds of formula I may also be delivered through the oral cavity by sublingual and/or buccal administration. Molded tablets, compressed tablets or freeze-dried tablets are exemplary forms that may be used. Exemplary compositions include those formulating the inventive compound(s) with fast-dissolving diluents such as mannitol, lactose, sucrose, and/or cyclodextrins. Also included in such formulations may be high molecular weight excipients such as celluloses (AVICEL®) or polyethylene glycols (PEG). Such formulations may also include an excipient to aid mucosal adhesion such as hydroxypropyl cellulose (HPC), hydroxypropyl methyl cellulose (HPMC), sodium carboxymethyl cellulose (SCMC), maleic anhydride copolymer (e.g., GANTREZ®), and agents to control release such as polyacrylic copolymer (e.g., CARBOPOL 934®). Lubricants, glidants, flavors, coloring agents and stabilizers may also be added for ease of fabrication and use.

Exemplary compositions for nasal aerosol or inhalation administration include solutions in saline which may contain, for example, benzyl alcohol or other suitable preservatives, absorption promoters to enhance absorption and/or bioavailability, and/or other solubilizing or dispersing agents such as those known in the art.

Exemplary compositions for parenteral administration include injectable solutions or suspensions which may contain, for example, suitable non-toxic, parenterally acceptable diluents or solvents, such as mannitol, 1,3-butanediol, water, Ringer's solution, an isotonic sodium chloride solution, 5 or other suitable dispersing or wetting and suspending agents, including synthetic mono- or diglycerides, and fatty acids, including oleic acid.

Exemplary compositions for rectal administration include suppositories which may contain, for example, suitable non-irritating excipients, such as cocoa butter, synthetic glyceride esters or polyethylene 10 glycols, which are solid at ordinary temperatures but liquefy and/or dissolve in the rectal cavity to release the drug.

Exemplary compositions for topical administration include a topical carrier such as PLASTIBASE® (mineral oil gelled with polyethylene).

The effective amount of a compound of the present invention may 15 be determined by one of ordinary skill in the art, and includes exemplary dosage amounts for an adult human of from about 0.05 to 100 mg/kg of body weight of active compound per day, which may be administered in a single dose or in the form of individual divided doses, such as from 1 to 4 times per day. It will be understood that the specific dose level and frequency of 20 dosage for any particular subject may be varied and will depend upon a variety of factors, including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the species, age, body weight, general health, sex and diet of the subject, the mode and time of administration, rate of excretion, drug combination, and severity of the 25 particular condition. Preferred subjects for treatment include animals, most preferably mammalian species such as humans, and domestic animals such as dogs, cats, horses, and the like, subject to cGMP-associated conditions.

The inventive compounds and compositions may be employed alone or in combination with each other and/or other suitable therapeutic 30 agents useful in treating cGMP-associated conditions such as other cGMP PDE inhibitors, particularly other cGMP PDE 5 inhibitors, modulators of the

large-conductance calcium-activated potassium (BK) channels, prostanooids, α -adrenergic agonists, endothelin antagonists, angiotensin II (especially, subtype AT₁) antagonists, angiotensin converting enzyme (ACE) inhibitors, renin inhibitors, and serotonin (5-HT_{2c}) agonists.

- 5 Exemplary of such other therapeutic agents are the following:
phenolamine, yohimbine, papaverine, apomorphine, sildenafil,
pyrazolopyrimidinones as described in U.S. Patent Nos. 5,272,147;
5,250,534; 5,426,107; and 5,346,901, quinazolinones as described in U.S.
Patent No. 5,482,941; AT₁ antagonists such as from losartan, irbesartan,
10 valsartan, and candesartan; ET_A antagonists such as bosentan, ABT-627,
and those described in U.S. Patent No. 5,612,359 and U.S. Patent
Application Serial No. 60/035,832, filed January 30, 1997; PDE 5 inhibitors
selected from imidazoquinazolines (see WO 98/08848), carbazoles (see WO
97/03675, WO 97/03985 and WO 95/19978), imidazopurinones (see WO
15 97/19947), benzimidazoles (see WO 97/24334), pyrazoloquinolines (see U.S.
Patent No. 5,488,055), quinazolinones as described in U.S. Patent No.
6,087,368, pyridines as described in U.S. Patent Application Serial No.
60/100,655 filed September 16, 1998, anthranilic acid derivatives (see WO
95/18097), fused heterocycles (see WO 98/07430) and thienopyrimidines
20 (see DE 19632423); and 5-HT_{2c} agonists selected from indoles (see *J. Med.
Chem.*, 40, 2762-2769 [1997], EP 655440 and EP 657426), and modulators
of the large-conductance calcium-activated potassium (BK) channels as
described in U.S. Patent Nos. 5,565,483 and 5,602,169, and in WO 98/04135
and WO98/23273.

- 25 The above other therapeutic agents, when employed in combination
with the compounds of the present invention, may be used, for example, in
those amounts indicated in the Physicians' Desk Reference (PDR) or as
otherwise determined by one of ordinary skill in the art.

- 30 The following assay can be employed in ascertaining the degree of
activity of a compound as a cGMP PDE inhibitor. Compounds described in

the following Examples have been tested in this assay, and have shown activity.

PDE Scintillation Proximity Assay Protocol

5 Sonicated human platelet homogenates are prepared by the method of Seiler, *et al.* (Seiler, S., Gillespie, E., Arnold, A.J., Brassard, C.L., Meanwell, N.A. and Fleming, J.S., "Imidazoquinoline Derivatives: Potent Inhibitors of Platelet Camp Phosphodiesterase which Elevate Camp Levels and Activate Protein Kinase in Platelets," *Thrombosis Research*, **62**: 31-42 (1991)). PDE 5 is abundant in human platelets, and accounts for approximately 90% of the cGMP hydrolytic activity in the homogenates. When necessary, PDE 5 can be resolved from other PDE activities in the homogenates by anion exchange chromatography on a fast protein liquid chromatography system (FPLC) using a Mono-Q anion exchange column (Pharmacia) eluted with a linear gradient of 10 mM - 450 mM NaCl.

15 The phosphodiesterase activity is assayed using a commercially available phosphodiesterase [^3H]cGMP scintillation proximity (SPA) assay kit (Amersham). The manufacturer's protocol is followed explicitly except that the reactions are carried out at RT and 3 mM nonradioactive cGMP is included in the suspension of SPA beads to prevent the synthesis of any additional radioactive products.

All documents cited in the present specification are incorporated herein by reference in their entirety.

25 The following Examples illustrate embodiments of the inventive compounds and starting materials, and are not intended to limit the scope of the claims. For ease of reference, the following abbreviations are used in the Examples, below:

Abbreviations

DMSO = dimethylsulfoxide

HPLC = high pressure liquid chromatography

30 LRMS = low resolution mass spectrometry

mp = melting point

tlc = thin layer chromatography

RT = room temperature

h = hour(s)

Ac = acetyl

5 Et = ethyl

Me = methyl

HOAc = acetate

EtOAc = ethyl acetate

EDAC • HCl = ethyl-3-(dimethylamino)propyl carbodiimide,

10 hydrochloride salt

HOBT = hydroxybenztriazole


NMP = N-methyl pyrrolidinone

TEA = triethylamine

15 **Preparation of Starting Materials**

Preparation 1


2-Amino-5-bromo-benzyl alcohol



20

To 13 mmol of methyl 2-amino-5-bromobenzoate in 20 mL of THF at RT was added 65 mmol of 1M lithium tri-tert-butoxyaluminumhydride over 10-15 minutes. The solution was heated at reflux for 17 hours. The solution was cooled, poured directly onto silica gel, and eluted with methylene chloride/ethyl acetate; 3:1 followed by 1:1. The product was eluted with 100% methanol to give 2.59g of the title compound as an off-white solid. MH⁺: 202; LC: 1.09'.


25

Preparation 2Diethyl 2-(4-cyanophenylamino)methylenemalonate

5

To 5.00 g (42.3 mmol) 4-aminobenzonitrile (Aldrich, 98%) was added 10.1 g (9.41 mL, 46.6 mmol, 1.1 eq) diethylethoxymethylenemalonate (Aldrich, 99%), and the mixture was dissolved in 50 mL toluene (HPLC grade). The solution was refluxed for four hours with a condensor open to the
10 air. The solution was then poured into 200 mL hexane, and the resulting white precipitate was filtered and washed well with more hexane to yield 10.62 g (37.0 mmol, 85% yield) of the title compound as a slightly off-white solid (per LC/MS and ¹H NMR). LC [MH⁺] 289, 97% purity.

15


Preparations 3-6Diethyl 2-(arylamino)methylenemalonate

20


Preparations 3-6 of formula (P1) having values for R₆, R₈ listed in Table 1 were prepared by the same method as in preparation 2, using the corresponding 4 amino-benzene.

TABLE 1

Preparation	R ₆	R ₈
3	Et	CN
4	Cl	CN
5	HOCH ₂	Br
6	CO ₂ CH ₃	Br

Preparation 7Diethyl 2-(2-acetoxymethyl-4-bromophenylamino)methylenemalonate


To a solution of 5.83 mmole of product from preparation 5 in 5ml of pyridine was added 8.74 mmole of acetic anhydride over 15 minutes and the mixture stirred at RT for 1.5 hours. Another 8.74 mmole of acetic anhydride were added, and the mixture stirred at RT for 15.5 hours. Another 2.94 mmole of acetic anhydride were added and the mixture stirred at RT for 4 hours. The mixture was poured into water, and an organic layer extracted with ethyl acetate. The organic layer was washed with water and saturated NaCl, dried over MgSO₄, and filtered. The filtrate was concentrated to give 2.10g of the title compound as an off-white solid. MH⁺: 416; LC: 4.25'.

Preparation 86-Cyano-4-hydroxyquinoline-3-carboxylic acid ethyl ester

- To 100 mL vigorously refluxing diphenyl ether (Aldrich) was added in open air, in portions over the course of one hour, 10.5 g (36.5 mmol) diethyl 2-(4-cyanophenylamino)methylenemalonate. The solution was refluxed for one additional hour. After allowing the solution to cool to below 100°C, it was poured into 200 mL hexane. The resulting precipitate was filtered and washed well with hexane to yield 8.39 g (34.6 mmol, 94%) of the title compound as a light brown solid: MH⁺: 243, mp >265°C.

Preparations 9-12

- 10 6,8-Disubstituted 4-hydroxyquinoline-3-carboxylic acid ethyl esters




(P2)

- 15 Preparations 9-12 of formula (P2) wherein R₆ and R₈ have the values listed in Table 2, were prepared by the same method as in preparation 8, using an appropriately substituted diethyl 2-(phenylamino) methylenemalonate.

20

TABLE 2

Preparation	R ₆	R ₈
9	CN	Et
10	CN	Cl
11	Br	AcOCH ₂ -
12	Br	-CO ₂ CH ₃


Preparation 134-Chloro-6-cyanoquinoline-3-carboxylic acid ethyl ester

5

To 9.50 g (39.2 mmol) ethyl 6-cyano-4-hydroxyquinoline-3-carboxylic acid was added 50 mL POCl_3 (Aldrich, 99%), and the resulting mixture was refluxed for 48 hrs. The POCl_3 was evaporated under reduced pressure, and the residue codistilled once with CHCl_3 , and twice with toluene.

10 The resulting brown solid was dissolved in CH_2Cl_2 and treated with triethylamine until aqueous washings of aliquots had $\text{pH} > 10$. The solution was then filtered through a 2" silica pad to yield 10.5 g (40.2 mmol, 103% yield) of the title compound as an off-white crystalline solid; MH^+ : 261, 97% purity.

15

Preparations 14-224-Chloroquinoline-3-carboxylic acid ethyl esters

20


The compounds of formula (P3) were prepared, wherein R_2 , R_6 , R_7 , and R_8 have the values listed in Table 3, using the same method as in preparation 13, with an appropriately-substituted ethyl 4-hydroxyquinoline-3-carboxylic acid.

25

TABLE 3

Preparation	R ₂	R ₆	R ₇	R ₈
14	H	CN	H	Et
15	H	CN	H	Cl
16	H	H	CF ₃	H
17	H	CF ₃	H	H
18	H	Br	H	AcOCH ₂
19	H	Br	H	CO ₂ CH ₃
20	Cl	Br	H	H
21	Cl	CN	H	H
22	Cl	CN	H	Et

5

Preparations 23 a) and b)a) Methyl 2-amino-5-cyanobenzoateb) Methyl 2-amino-3-ethyl-5-cyanobenzoate

10

, wherein R₈ is H or Et


15 A mixture of methyl 2-amino-5-bromobenzoate (4.6 g, 20 mmol),
 and CuCN (1.97 g, 22 mmol) in NMP (20 mL) was heated to 190°C and
 stirred for 3 h. The reaction mixture was poured into a solution of ethylene
 diamine (4 mL) in H₂O (16 mL) and extracted with toluene (4x20 mL). The
 combined extracts were dried (Na₂SO₄). Removal of the solvent gave the title
 compound a) (*i.e.*, R₈ is hydrogen). Use of methyl 2-amino-5-bromo-3-ethyl
 20 bromobenzoate gave the title compound b) (*i.e.*, R₈ is ethyl).

Preparations 24 a) and b)

a) 6-Bromo-2,4-dihydroxyquinoline-3-carboxylic acid ethyl ester


b) 6-Cyano-2,4-dihydroxyquinoline-3-carboxylic acid ethyl ester

5

wherein R₆ is Br or CN.

- Diethyl malonate was added to a freshly prepared solution of sodium ethoxide in ethyl alcohol. The resulting mixture was stirred for 30 min. To it was then added dropwise a solution of 2-amino-5-bromobenzoic acid methyl ester in ethyl alcohol. The resultant mixture was refluxed overnight. The ethyl alcohol was removed under reduced pressure, and the residue was dissolved in water and extracted with EtOAc. The aqueous layer was acidified with glacial AcOH to pH5. The precipitate was collected by filtration to yield 2.6 g, (94% yield) the title compound a) (wherein R₆ is Br). Use of 1.76 g (10 mmol) 2-amino-5-cyanobenzoic acid methyl ester yielded 1.7 g (66% yield) of the title compound b) (wherein R₆ is CN).

20

Preparation 256-Cyano-2,4-dihydroxy-8-ethylquinoline-3-carboxylic acid ethyl ester


25

To a solution of methyl 2-amino-5-cyano-3-ethylbenzoate (1.36 g, 6.2 mmol) and TEA (2.6 mL, 18.6 mmol) in THF (50 mL) was added methyl malonyl chloride (1 mL, 9.3 mmol). The mixture was stirred at RT for


overnight. Additional TEA (2.6 mL) and methyl malonyl chloride (1 mL) were added, and the reaction was continued for another 24 h. The mixture was diluted with EtOAc and washed with H₂O and brine. The solvent was removed, and the residue was chromatographed (silica gel, EtOAc/hexane, 3:7) to give the amide intermediate. NaH (400 mg, 60% oil dispersion) was washed with dry hexane and dried under nitrogen. To it was added a solution of methyl 2-amino-5-cyano-3-ethylbenzoate (1 g, 3 mmol) in toluene (40 mL), which was followed by dropwise addition of EtOH (4 mL). The resultant mixture was refluxed for overnight. The solvent was removed and the residue was dissolved in H₂O (insoluble particulates were removed by filtration). Upon acidification with HOAc, the title compound (590 mg) was precipitated out and collected by filtration.

Preparation 26

6-Bromo-2-(2-pyridinyl)quinoline-4-carboxylic acid



To a suspension of 5-bromo isatin (5 g, 20.5 mmol) and 2-acetylpyridine (2.5 g, 20.5 mmol) in aqueous KOH (1.2 g, 33%) was added EtOH until a solution formed. The mixture was refluxed over night. The reaction mixture was neutralized with HOAc (50%) and filtered. The solid was resuspended in EtOH and filtered to give the title compound (5.5 g). LC: 1.66 and MH⁺: 329.


Preparation 27Azido-6-bromo-2-(2-pyridinyl)quinoline-4-carboxylate

5


A mixture of 6-bromo-2-(2-pyridine)quinoline-3-carboxylic acid (2.5 g) in SOCl_2 (30 mL) was refluxed for 2 h, and the excess SOCl_2 was removed. The residue was resuspended in acetone (100 mL). To the suspension was added a solution of NaN_3 (0.5 g) in water (50 mL) and stirred
10 for 2h. The solid product was collected by filtration and washed thoroughly with water, then dried at 40°C overnight to give the title compound (2.3 g).
LC: 2.13'.

Preparation 28

15

6-Bromo-4-*t*-butoxycarbonylamino-2-(2-pyridinyl)quinoline

A mixture of preparation 27 (2 g) in toluene (50 mL) was refluxed for
20 3h. Then *t*BuOH (1 mL) was added and continue to reflux for 2 h. The reaction mixture was cooled to RT and filtered. The filtrate was concentrated to give the title compound (2 g).

Example 14-[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyanoquinoline-3-carboxylic acid ethyl ester


5

To 10.0 g (38.4 mmol) 4-chloro-6-cyanoquinoline-3-carboxylic acid ethyl ester was added 10.4 g (50 mmol, 1.5 eq) 3-chloro-4-methoxybenzyl amine hydrochloride and 35 mL (228 mmol, 6 eq) diisopropylethylamine (Aldrich, 99.5+%). The mixture was dispersed in 200 mL *n*-propanol and brought to reflux for two hours. The solution was then poured into 500 mL water. The resulting precipitate was filtered, then washed with 500 mL water, then twice washed with 50 mL absolute ethanol to yield 14.87 g (37.6 mmol, 97% yield) of the title compound as a white fibrous solid after codistilling with absolute ethanol: mp: 161-162°C; LC/MS (M/Z) 396 [M+H] observed, 97% purity.

15











Examples 2-554-Aminoquinoline-3-carboxylic acid ethyl esters















20
































Using the same or similar method of Example 1, compounds having the formula (Iae) were prepared, wherein R_2 , R_6 , R_7 , and R_8 have the values listed in Table 4, starting with corresponding 3-carboxylate-4-chloroquinolines and amine hydrochlorides.


TABLE 4

Ex. No.	$-R_5$	R_2	R_6	R_7	R_8	HPLC retention time (min.)	MH^+
2		H	H	CF_3	H	3.63	439
3		Cl	Br	H	H	4.75	437
4		Cl	Br	H	H	4.56	485
5		H	CF_3	H	H	3.64	439
6		H	CN	H	H	3.73	352
7		H	CN	H	H	3.82	352
8		H	F	H	H	3.41	389
9		H	Br	H	CO_2nPr	3.29	535
10		H	Br	H	CO_2Me	3.52	507
11		H	Br	H	H	2.82	449

12		H	CN	H	Et	2.78	424
13		H	CN	H	Cl	3.89 ^b	430
14		H	Br	H	AcOCH ₂	1.73 ^a	521
15		H	Br	H	Me	3.68	463
16		H	CN	H	Et	1.56 ^a	394
17		H	CN	H	Et	1.68 ^a	428
18		H	CN	H	Et	1.65 ^a	428
19		H	CN	H	Et	1.46 ^a	390
20		H	CN	H	Et	1.73 ^a	380
21		H	CN	H	Et	1.73 ^a	380
22		H	CN	H	CO ₂ nPr	3.43	482
23		H	CN	H	CO ₂ Me	3.11	454
24		H	CN	H	CO ₂ Et	3.27	468
25		H	CN	H	Et	1.23 ^a	350

26		H	CN	H	Et	1.33 ^a	404
27		H	CN	H	Et	1.49 ^a	428
28		H	CN	H	Et	1.38 ^a	396
29		H	CN	H	Et	1.49 ^a	428
30		H	CN	H	Et	1.65 ^a	402
31		H	CN	H	Et	1.70 ^a	416
32		H	CN	H	Et	1.60 ^a	428
33		H	CN	H	Et	1.42 ^a	420
34		H	CN	H	Et	1.39 ^a	420
35		H	CN	H	Et	1.53 ^a	444
36		H	CN	H	Et	1.53 ^a	442
37		H	CN	H	Et	1.28 ^a	550
38		H	CN	H	Et	1.55 ^a	446


39		H	CN	H	Et	1.34 ^a	444
40		H	CN	H	Et	3.54	374
41		H	CN	H	Et	3.55	374
42		H	CN	H	Et	3.94	452
43		H	CN	H	Et	3.95	452
44		H	CN	H	Et	3.78	388
45		H	CN	H	Et	3.77	388
46		H	CN	H	Et	3.26	360
47		H	CN	H	Et	3.65	438
48		H	CN	H	Et	3.53	374
49		H	CN	H	Et	1.38 ^a	426
50		H	CN	H	Et	1.46 ^a	406
51		H	CN	H	Et	1.46 ^a	386
52		H	CN	H	Et	1.80 ^a	442
53		H	CN	H	Et	1.41 ^a	434
54		H	CN	H	Et	1.36 ^a	448

55		H	CN	H	Et	1.52 ^a	424
----	---	---	----	---	----	-------------------	-----


Example 56

6-Bromo-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-hydroxymethylquinoline-3-carboxylic acid ethyl ester

5




A mixture of 1.84 mmole 6-bromo-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-8-(acetoxymethyl)quinoline-3-carboxylic acid ethyl ester (*i.e.*, Example 14) and 1.84 mmole of potassium carbonate in 20 mL of ethanol was stirred at RT for 20 hours. The mixture was concentrated, and the residue was diluted with water and extracted with ethyl acetate. The resultant organic layer was dried over MgSO₄ and filtered, and the filtrate concentrated to give 0.85g of the title compound as a tan solid. MS: 481; LC: 3.43'.

Example 576-Bromo-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-(chloromethyl)quinoline-3-carboxylic acid ethyl ester

5


A solution of 1.73 mmole 6-bromo-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino)-8-hydroxymethylquinoline-3-carboxylic acid ethyl ester (*i.e.*, Example 56) in 8 ml of thionyl chloride was stirred at RT for 1.5 hours. The solution was concentrated to give 0.86 g of the title compound as an orange foam. MS: 499; LC: 4.15'

10

Example 586-Bromo-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-(*N,N*-dimethylaminomethyl)quinoline-3-carboxylic acid ethyl ester

5

A solution of 0.20 mmole of example 57 and 2 mmole of 2M dimethylamine in 1ml of THF was stirred at RT for 22 hours. An additional 0.25 ml of 2M dimethylamine in THF was added and the solution stirred for 4
10 hours. The solution was diluted with ethyl acetate and washed with water. The organic layer was dried over MgSO_4 and filtered, and the filtrate was concentrated. The residue was chromatographed on silica gel eluted with methylene chloride followed by methylene chloride/ethyl acetate; 95:5 and finally with ethyl acetate to give 71 mg of the title compound as a beige solid.
15 MS: 507; LC: 3.71'

Example 594-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinecarboxylic acid

5


To 8.0 g (20 mmol) 4-(3-Chloro-4-methoxyphenylmethylamino)-6-cyanoquinoline-3-carboxylic acid ethyl ester was added 100 mL THF, 100 mL MeOH and 100 mL 1M NaOH, and the resulting mixture was stirred well. The solids gradually dissolved as the reaction progressed. After 1h, the THF and most of the MeOH was evaporated under reduced pressure to leave an aqueous slurry of a white solid. The slurry was acidified to pH 1.5 with HCl, stirred well for 1h, and then filtered and washed well with water to give 7.5 g (20 mmol, ~100% yield) of a white chalky solid. Codistilling twice to dryness with absolute ethanol afforded the title compound: LC/MS: M/Z 368 (M+H) observed, 90+% purity; mp: decomposed 255-260°C. LC 2.93'.

10

15

Examples 60-64

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-quinolinecarboxylic acids




5

Using the same or similar method as described in Example 59, compounds having the formula (Iaf) were prepared, wherein R_2 , R_6 , R_7 , and R_8 have the values listed in Table 5, starting with corresponding 3-carboxylatequinolines.

10


TABLE 5

Ex. No.	R_2	R_6	R_7	R_8	HPLC retention time (min.)	MH^+
60	H	H	CF_3	H	3.67	411
61	Cl	Br	H	H	4.29	455
62	H	F	H	H	3.31	361
63	H	CN	H	Et	2.56	396
64	H	CN	H	Cl	3.23	402

Example 654-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-quinolinecarbonitrile

5


To 2.16 g (5.87 mmol) 4-[[[(3-chloro-4-methoxyphenyl)-
methyl]amino]-6-cyano-3-quinolinecarboxylic acid was added 50 mL diphenyl
ether (Aldrich), and nitrogen was vigorously bubbled through this mixture for
20 minutes. The resulting mixture was then heated to 240 °C under nitrogen
10 for 30 minutes. During the course of the reaction, the dispersed starting
material slowly dissolved, and gas evolution was observed. The resulting
solution was poured into 200 mL hexane, and a precipitated solid was filtered
to yield 1.43 g (4.42 mmol, 75% yield) of the title compound as a off-white
solid: LC/MS: M/Z 324 (M+H) observed, 96% purity; mp: decomposed 227-
15 230 °C. LC: 2.84'.

Example 664-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-6-quinolinecarbonitrile

5

The reaction procedure used was similar to that described for Example 65 with the following changes: 101 mg (0.255 mmol) of 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-cyano-8-ethyl-3-quinolinecarboxylic acid was used to afford 51 mg (57% yield) of the title compound. LC: 1.20' M/Z

10 352 mp: 206-207 °C.

Example 674-[[[(3-Chloro-4-methoxyphenyl)methyl]N-methylamino]-6-quinolinecarbonitrile

15

To 250 mg 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-quinolinecarbonitrile (*i.e.*, Example 65) was added 50 mg of NaH, and the mixture was flushed with nitrogen. 5 mL of DMF was added, and it was stirred for half hour. To the mixture was then added 70 μ L of CH₃I, and it was


20

stirred at RT. After 1 hour the reaction was quenched with aqueous NH_4Cl and diluted with water, and the precipitate collected by filtration to give (153 mg) of the title compound. LC: 2.48; M/Z 338 m.p. 153-155°C.

5

Example 68

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyanoquinoline-3-carboxylic acid pentafluorophenyl ester




10

To 275 mg (0.748 mmol) of 4-[[[(3-chloro-4-methoxyphenyl)-methyl]amino]-6-cyano-3-quinolinecarboxylic acid was added 276 mg (1.50 mmol, 2.0 eq) pentafluorophenol, and the mixture was dispersed in 5 mL anhydrous DMF under nitrogen. To this dispersion was added a solution of 231 mg (1.12 mmol, 1.5 eq) of dicyclohexylcarbodiimide (Aldrich) in 2 mL anhydrous ethyl acetate at RT. The resulting mixture was stirred for 18 hours. The reaction was then diluted with 10 mL EtAc, and the white precipitate was filtered and discarded. The filtrate was concentrated *in vacuo*, and the residue was triturated with 5% MeOH/95% CH_2Cl_2 to give 164 mg (0.307 mmol, 41% yield) of the title compound as a white solid: LC/MS: M/Z 534 (M+H) observed, 100% purity; mp: decomposed 215-217 °C. LC: 4.15.

20

Example 69

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-8-ethylquinoline-3-
carboxylic acid pentafluorophenyl ester




5

The reaction procedure used was similar to that described for
Example 68 with the following changes: 3.95 g (10.0 mmol) 4-[[[(3-chloro-4-
methoxyphenyl)methyl]amino]-6-cyano-8-ethyl-3-quinolinecarboxylic acid was
10 used to afford 2.81 g (5.00 mmol, 50% yield) of the title compound as a white
solid: mp: 169-170 °C; LC: 2.14^a; MH⁺ 562.

Examples 70-122

15 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-quinolinecarboxamides









Compounds having the formula (Ia), wherein R_2 , R_6 , R_7 , R_8 , R_{10} , and R_{11} , have the values listed in Table 6, were prepared by the following method, starting with corresponding 3-carboxylatequinolines and an appropriate amine:
















5

















To 1 eq of 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-quinolinecarboxylic acid pentafluorophenyl ester in THF (approx. 25 mg per 1 mL THF) was added 2 eq of an appropriate primary or secondary amine. The resulting reaction solution was allowed to stir for 18 hours at RT. The THF was then evaporated *in vacuo*, and the residue was triturated with a 1:1 solution of ether/hexane. The precipitated solid was filtered and washed with 1:1 ether/hexane to yield the desired 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-quinolinecarboxamide. Yields were generally > 25%.


15

TABLE 6

Ex. No.	R_2	R_6	R_7	R_8	$-NR_{10}R_{11}$	HPLC retention time (min.)	MH^+
70	H	H	CF_3	H	$-NH_2$	3.35	410
71	H	H	CF_3	H		3.09	501
72	H	H	CF_3	H		3.06	515
73	H	H	CF_3	H		1.78 ^a	523
74	H	H	CF_3	H		1.81 ^a	482
75	H	H	CF_3	H		1.92 ^a	508
76	H	H	CF_3	H		1.92 ^a	438

77	H	H	CF ₃	H		1.73 ^a	494
78	H	CN	H	H		2.49	458
79	H	CN	H	H		2.56	480
80	H	CN	H	H		2.45	507
81	H	CN	H	H		2.18	450
82	H	CN	H	H	-NH ₂	2.73	367
83	H	CN	H	H		2.99	423
84	H	CN	H	H		1.57 ^a	438
85	H	CN	H	H		2.57	410
86	H	CN	H	H		1.47 ^a	503
87	H	CN	H	H		1.91 ^a	478
88	H	CN	H	H		1.55	464
89	H	CN	H	H		1.80	504
90	H	CN	H	H		1.63	494
91	H	CN	H	H		1.64	504
92	H	CN	H	H		1.70	518
93	H	CN	H	Et	-NH ₂	2.34	395


94	H	CN	H	Et		2.20	486
95	H	CN	H	Et		2.05	532
96	H	CN	H	Et		1.92	506
97	H	CN	H	H		2.38	478
98	Cl	Br	H	H		3.36	551
99	Cl	Br	H	H		3.38	565
100	Cl	Br	H	H		3.48	591
101	Cl	Br	H	H		3.49	545
102	Cl	Br	H	H		2.74	537
103	Cl	Br	H	H		1.57 ^a	539
104	Cl	Br	H	H		1.52 ^a	594
105	Cl	Br	H	H		1.55 ^a	581
106	Cl	Br	H	H		3.61	579
107	Cl	CN	H	H		3.09	492
108	Cl	CN	H	Et		1.73 ^a	520
109	H	CN	H	Et		1.35 ^a	449

110	H	CN	H	Et		1.21 ^a	532
-----	---	----	---	----	---	-------------------	-----

Examples 111-179

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinecarboxamides

5

































10 Compounds having the formula (Iah), wherein R_{10} and R_{11} have the values listed in Table 7, were prepared by the following method:

















To a solution of 1.0 eq of the appropriate primary amine or 2.0 eq of the appropriate secondary amine in anhydrous THF (3 mL) was added 15 mg 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinecarboxylic acid pentafluorophenyl ester (0.028 mmol). Where the amine was a
 15 hydrohalide salt, 3 eq of triethylamine were also added. The solutions were stirred overnight at 50 °C, then filtered and purified by SCX cartridges (2 g capacity, *p*-toluene sulfonic acid) on a Bohdan SPD robot (10 mL MeOH wash, 10 mL 2M NH_3 /MeOH elution). The desired 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinecarboxamide was
 20 recovered either by filtration (manual), by concentration of the SCX eluent (by Savant Speedvac), or (if further purification was required) by preparative HPLC (Shimadzu). Average purity: 93% (LC/MS). Average yield: > 25%.

TABLE 8

Ex. No.	$-NR_{10}R_{11}$	HPLC retention time (min.) ^a	MH ⁺
111		1.63	395
112		1.83	423
113		1.96	435
114		2.17	451
115		1.84	439
116		1.70	455
117		1.81	439
118		2.01	467
119		1.97	451
120		1.69	411
121		1.66	425
122		0.64	439
123		1.77	453
124		2.16	453
125		1.78	453
126		1.57	455

127		1.70	465
128		1.78	453
129		1.83	467
130		1.66	452
131		1.77	492
132		1.59	466
133		1.49	466
134		1.49	480
135		1.74	494
136		1.60	464
137		1.66	478
138		1.48	492
139		1.53	424
140		1.53	440
141		1.58	452

142		1.54	452
143		1.58	480
144		1.52	494
145		1.63	506
146		1.59	478
147		1.46	478
148		1.96	522
149		1.73	540
150		1.75	514
151		1.54	466
152		1.63	494
153		1.63	437
154		1.78	453
155		1.66	452
156		1.55	494


157		1.59	437
158		1.59	478
159		1.43	464
160		1.83	435
161		1.59	451
162		1.69	479
163		1.91	507
164		1.63	478
165		1.63	478
166		1.90	493
167		1.43	436
168		1.43	521
169		1.41	480
170		1.64	518
171		1.48	513
172		1.76	514

173		1.45	450
174		1.44	464
175		2.00	457
176		1.61	458
177		1.87	458
178		1.86	487
179		2.25	501

Example 180

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-(hydroxymethyl)-6-quinolinecarbonitrile

5




To 530 mg (1.0 mmol) of 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinecarboxylic acid
 10 pentafluorophenyl ester was added 15 mL of anhydrous DMF, and the resulting mixture was cooled to 0 °C. A solution of 75 mg of NaBH₄ in 5 mL of anhydrous DMF was added to the reaction mixture, resulting in a clear red

solution with gas evolution. After stirring for 1 hour at 0 °C, the reaction was quenched with 2% TFA/MeOH, and the solvent was removed under reduced pressure. The oily residue was purified by flash chromatography (silica gel, 3-5% MeOH/CH₂Cl₂) to afford 87 mg (25% yield) of the title compound. LC: 5 2.60'; MH⁺: 354.


Example 181

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-(hydroxymethyl)-6-quinolinecarbonitrile



To a suspension of 100 mg (0.236 mmol) of 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-cyano-8-ethylquinoline-3-carboxylic acid ethyl ester in 2.8 mL anhydrous THF under nitrogen, was added 1.2 mL (5.0 eq) of 1M lithium tri-*t*-butoxyaluminumhydride/THF. The resulting mixture was refluxed overnight with stirring. The reaction was quenched with 1 mL of MeOH and partitioned between 75 mL 1M NaOH and 100 mL CH₂Cl₂. The organic phase was washed with 1M NaOH (2x30 mL) and dried over MgSO₄. Removal of the solvent under reduced pressure gave 75 mg (84% yield) of the title compound as a light yellow solid. The analytical sample was obtained by trituration in ether: mp: 188-189 °C. LC: 3.15'; MH⁺: 382.

Examples 182-190

4-Amino-3-(hydroxymethyl)quinolines

(Iai)

5

Using the same method of Example 181, compounds having the formula (Iai) were prepared, wherein R_5 , R_6 , and R_8 have the values listed in Table 8, using an appropriate amine (having the group R_5). In Example 190,

10 the reaction was started with 3-ethyl-8-methyldicarboxylatequinoline.

TABLE 8


Ex. No.	$-R_5$	R_6	R_8	HPLC retention time (min.)	MH^+ Found
182		Br	H	2.41	407
183		CN	Cl	2.92	388
184		CN	Et	1.50 ^a	338
185		CN	Et	1.51 ^a	338
186		CN	Et	1.45 ^a	386
187		CN	Et	1.43 ^a	386

188		CN	Et	1.38 ^a	352
189		CN	Et	1.25 ^a	348
190		CN	-(CH ₂)OH	2.59	384

Examples 191-194

3-Aminomethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-quinolinecarbonitriles

5






- 10 Compounds having the formula (Iaj) were prepared, wherein R₁₀ and R₁₁ have the values listed in Table 9, by the following method.

- To 70 mg 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-(hydroxymethyl)-6-quinolinecarbonitrile was added 5 mL anhydrous CH₂Cl₂ and 70 μL (0.44 mmol, 2.0 eq) diisopropylethylamine, and the resulting
 15 solution was cooled to 0°C under nitrogen. Methanesulfonyl chloride (17 μL, 0.22 mmol, 1.1 eq) was then added, and the resulting reaction solution was stirred at 0°C for 30 minutes. The solution was then quenched with 0.24 mmol (1.1 eq) of the appropriate amine (having groups R₁₀, R₁₁), and the resulting solution was stirred for 30 minutes. The solvent was evaporated

under reduced pressure and the residue was chromatographed in 5% MeOH/CH₂Cl₂ to yield the title compound.


TABLE 9

Ex. No.	-NR ₁₀ R ₁₂	HPLC retention time (min.)	MH ⁺
191		2.39	436
192	-NH ₂	1.12	353
193		1.83	444
194		1.84	421

5

Example 195

6-Bromo-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-
[(4-methyl-1-piperazinyl)methyl]] quinoline



10

To 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-(hydroxymethyl)-6-bromoquinoline was added SOCl₂, and the resulting reaction solution was stirred at RT under nitrogen for 30 minutes. The SOCl₂ was then evaporated under reduced pressure, and the residue was left overnight under high vacuum. The yellow residue was redissolved in anhydrous DMF, and this solution was then added to a solution of N-methylpiperazine in anhydrous


15

THF. The resulting reaction solution was stirred for 6 hours at RT, by which time HPLC showed the reaction to be complete. After evaporating the solvent, the crude product was chromatographed in 3% MeOH/CH₂Cl₂ to yield the title compound. LC: 2.12'; MH⁺: 489

5

Examples 196-224

3-Aminomethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-quinolinecarbonitriles

















10














Compounds having the formula (Iak) were prepared, wherein values for R₂, R₈, R₁₀ and R₁₁ are as in Table 10, using the same or similar method as in Example 195, starting with the corresponding 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-(hydroxymethyl)-6-quinolinecarbonitrile.

15

TABLE 10


Ex. No.	-NR ₁₀ R ₁₁	R ₂	R ₈	HPLC retention time (min.)	MH ⁺ Found
196		H	H	2.55	437
197		H	H	3.43	465

198		Cl	H	1.43 ^a	471
199		H	H	2.40	381
200		H	H	2.43	449
201		H	H	1.96	437
202		H	H	2.21	435
203		H	H	2.24	451
204		H	H	1.81	437
205		H	H	1.60	423
206		H	H	1.97	451
207		H	H	2.33	435
208		H	H	2.61	435
209		H	H	1.86	423
210		H	H	2.74	449
211		H	H	2.25	451

212		H	H	1.17 ^a	437
213		H	H	2.80	477
214		H	H	2.50	461
215		H	H	2.45	473
216		H	Et	1.34 ^a	465
217		H	H	2.40	443
218		H	Et	1.27 ^a	465
219		H	Cl	1.20 ^a	457
220		H	Cl	1.23 ^a	485
221		H	Cl	1.35 ^a	457
222		H	Et	1.20 ^a	451
223		H	Et	1.45 ^a	479
224		H	Et	1.45 ^a	479

Examples 225-263

3-Aminomethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-
quinolinecarbonitriles



5

Compounds of formula **(Ia)** were prepared, wherein R_{10} and R_{11} have the values listed in Table 10, with the following method (Automated parallel synthesis).



















10


















A mixture of 1 eq. of 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-chloromethyl-6-cyanoquinoline and 3 eq. of an appropriate amine in THF was reacted for overnight at RT. The reaction mixture was filtered and the filtrate was concentrated. The residue was dissolved in CH_2Cl_2 and washed with H_2O , then dried. Removal of the solvent gives the product.

15

TABLE 11

Ex. No.	$-NR_{10}R_{11}$	HPLC retention time (min.)	MH ⁺
225		2.54	407
226		2.65	421
227		2.82	435
228		2.99	420


229		2.42	411
230		2.83	438
231		2.4	393
232		2.46	407
233		2.92	423
234		3.18	438
235		2.76	435
236		2.48	395
237		2.42	411
238		2.99	437
239		2.73	423
240		2.8	423
241		3.28	435
242		2.84	423
243		2.44	381
244		2.5	410
245		2.4	397
246		3.07	437

247		2.91	423
248		2.55	395
249		2.44	395
250		2.55	409
251		2.64	409
252		2.56	406
253		3	423
254		2.65	425
255		2.54	409
256		3.09	437
257		2.44	411
258		2.43	410
259		2.61	409
260		2.61	409
261		2.64	437
262		2.71	425
263		2.63	437

Examples 264-266

3-(*N*-acylaminomethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-quinolinecarbonitriles


5



Compounds of formula **(Iam)** were prepared, wherein R_{12} has the values in Table 12, below. To a mixture of acyl chloride and triethylamine in CH_2Cl_2 was added a solution of 3-aminomethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-6-quinolinecarbonitrile (*i.e.*, Example 192) in dioxane. The reaction mixture was stirred at RT for 1 h. Then the mixture was diluted with EtOAc and washed with water brine and dried (MsO_4). The solvent was removed under reduced pressure and the residue was subjected to flash column chromatography (silica gel, $CH_2Cl_2/MeOH$, 10:1) to give the compounds of Table 12.


TABLE 12

Ex. No.	R_{12}	HPLC retention time (min.)	MH ⁺
264		2.76	457
265		3.00	463

266		2.50	458
-----	---	------	-----

Examples 267-283












5 3-Alkoxyethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]quinolines






Compounds of formula **(Ia)** were prepared, wherein R₂, R₆, R₈, and R₉ have the values listed in Table 13, with the following method.

To 4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-3-(hydroxymethyl)-6-bromoquinoline was added SOCl₂, and the resulting reaction solution was stirred at RT under nitrogen for 30 minutes. The SOCl₂ was then evaporated under reduced pressure, and the residue was left overnight under high vacuum. The yellow residue was redissolved in anhydrous DMF, and this solution was then added to a solution of appropriate alcohol in anhydrous THF (or the yellow residue was treated directly with an appropriate liquid alcohol). The resulting reaction solution was stirred for 6 hours at elevated temperature, by which time HPLC showed the reaction to be complete. After evaporating the solvent, the crude product was chromatographed to yield the title compound.

TABLE 13


Ex. No.	-R ₉	R ₂	R ₆	R ₈	HPLC retention time (min.)	MH ⁺
267		H	CN	Et	1.69 ^a	424
268	-CH ₃	H	CN	Et	3.10	396
269		H	CN	Et	3.35	466
270		H	CN	Et	3.42	424
271		H	CN	Et	3.29	410
272		H	CN	Et	1.60 ^a	454
273		H	CN	Et	1.39 ^a	426
274		H	CN	Et	3.91	565
275		H	CN	Et	3.91	565
276		H	CN	Et	3.80	551
277		H	Br	H	2.78	435
278		Cl	CN	H	2.13 ^a	571

279		Cl	CN	Et	2.37 ^a	599
280	CH ₃	H	Br	ClC H ₂	3.55	469
281		H	Br	ClC H ₂	3.78	497
282		H	CN	Et	3.18	438
283	CH ₃	H	CN	ClC H ₂	3.17	439

Examples 284-288

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[[[(2S)-2-pyrrolidiny]]methoxy]methyl]-6-quinolinecarbonitriles

5



Compounds of formula **(IaO)** were prepared, wherein R₂ and X₁ have the values listed in Table 14, with the methods set forth below.

10

Example 284: Example 275 was treated with TFA/CH₂Cl₂ (1:1) for 2 hours.

The reaction mixture was concentrated. The residue was treated with NaHCO₃ (aq. 10%) and extracted with CH₂Cl₂. The combined extracts were dried (NaSO₄) and the solvent was removed. The residue was then purified by preparative HPLC to give Example 284.

15

Example 285: The same method as Example 285 was used, starting with Example 279.


- 5 Example 286: To a solution of Example 284 (30 mg) in THF was added triethylamine (28 μ L) which was followed by CH_3I (4.5 μ L). The resultant mixture was stirred at RT for overnight. The solvent was removed and the residue was chromatographed (silica gel, $\text{CH}_3\text{OH}/\text{CHCl}_3$, 1:9) to give Example 286 (5.1 mg).

10

- Examples 287 & 288: To a solution of 0.127 mmole of Example 284 and 1.27 mmole of the ketone in 0.5 mL of methanol cooled in an ice bath was added 0.635 mmole of sodium cyanoborohydride. The ice bath was removed and the suspension stirred for 4 hours to 2 days. The solution was acidified with concentrated HCl and allowed to stir for a few minutes. The solution was
- 15 neutralized with concentrated NH_4OH and extracted with methylene chloride. The organic layer was dried over MgSO_4 and filtered, and the filtrate concentrated. The residue was chromatographed on silica gel eluted with methylene chloride/methanol; 95:5 and 9:1 to give Examples 287 and 288.


20

TABLE 14

Ex. No.	R_2	X_1	HPLC retention time (min)	MH^+
284	H	H	1.27 ^a	465
285	Cl	H	1.83 ^a	499
286	H	$-\text{CH}_3$	1.27 ^a	479
287	H		2.72	533
288	H	$-\text{CH}(\text{CH}_3)_2$	2.57	507

Examples 289-296

8-Aminomethyl-3-alkoxymethyl-4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]quinolines









5

Compounds having the formula **(Iap)** were prepared, wherein R_8 , R_9 , R_{13} and R_{14} have the values listed in Table 15. A solution of 0.072 mmole of 8-chloromethyl quinoline and 0.072 mmole of amine in 0.5 mL of THF was stirred at RT for 16 hours. Another 0.144 mmole of amine were added to the incomplected reactions and the solution stirred for 24 hours. The solution was diluted with water and extracted with ethyl acetate. The organic layer was washed with water, dried over $MgSO_4$, and filtered, and the filtrate was concentrated. The residue was chromatographed on silica gel eluted with methylene chloride/methanol; 98:2, 95:5 followed by methylene chloride/methanol/ NH_4OH ; 90:9:1, to give the products.

15

TABLE 15


Ex. #	$-R_9$	R_8	$-NR_{13}R_{14}$	HPLC retention time (min.)	MH^+
289	$-CH(CH_3)_2$	Br	$\begin{array}{c} CH_3 \\ \\ -N- \\ \\ CH_3 \end{array}$	1.54 ^a	506
290	$-CH(CH_3)_2$	Br	$-NHCH_3$	1.47 ^a	492

291	-CH ₃	CN		2.81	425
292	-CH ₃	CN		2.15	480
293	-CH ₃	CN		1.60	468
294	-CH ₃	CN		2.30	508
295	-CH ₃	CN		1.62	494
296	-CH ₃	CN		1.62	508

Examples 297-299

3-Alkoxyethyl-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-6-quinolinecarbonitriles




5



Compounds having the formula (Iaq) were prepared, wherein R₉ is substituted alkyl comprising CH₂CH(OH)CH₂NR₁₀R₁₁, and R₁₀ and R₁₁ have the values listed in Table 16. A solution of 0.11 mmole of Example 282 and 0.22 mmole of the appropriate amine in 1 mL of propanol was heated at 60-70°C for 45 minutes to 30 hours. The solution was concentrated and the

residue chromatographed (silica gel, methylene chloride/methanol; 95:5, 9:1 and 4:1) to give the products.

TABLE 16


Ex. No.	-NR ₁₀ R ₁₁	HPLC retention time (min.)	MH ⁺
297		2.51	509
298		2.67	539
299		2.48	483

5

Example 300

3-[[[3-Bis(1-methylethyl)amino]-2-methoxypropoxy]methyl]-4-[[[(3-chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-6-quinolinecarbonitrile]

10




To 0.069 mmole of Example 298 in 0.5ml of DMF cooled in an ice bath was added 0.072 mmole of 60% NaH in mineral oil. The ice bath was removed, the suspension stirred for 90 minutes, and 0.076 mmole of iodomethane were added. The solution was stirred at RT for 16 hours. An

15

additional 0.038 mmole of 60% NaH in mineral oil and 0.038 mmole of iodomethane were added and the solution stirred for 4 hours. The solution was diluted with water and extracted with ethyl acetate. The organic layer was washed with water and saturated NaCl solution, dried over MgSO₄ and filtered, and the filtrate concentrated. The residue was chromatographed on silica gel eluted with methylene chloride/methanol; 95:5 to give 19 mg of the title compound as a colorless oil. MH⁺: 553; LC: 3.21'.

Examples 301 and 302

10 [[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinyl]methyl]propanedioic acid dialkyl ester



15 Compounds having the formula (Iar) were prepared, wherein R₈ and R₉ have the values listed in Table 17. To a dry clean reaction flask was charged 1.3 g (33 mmole) of sodium hydride (60% in mineral oil) and 160 ml of dry THF. To this mixture was added dropwise 6.5 g (30 mmole) of di-*t*-butylmalonate over a period of 20 minutes at RT. After completion of the addition, the reaction mixture was stirred at RT for an additional 15 minutes then cooled to -78°C. A solution of the 3-chloromethyl quinoline (~10 mmole) in 160 ml of DMF was added *via* an addition funnel over a period of 1 hour. After completion of the addition, the reaction mixture was stirred at -78°C for one more hour then was quenched with water at 0°C. THF was removed under reduced pressure. The resulting solution was diluted with

25

ethyl ether washed with water three times, brine, dried over sodium sulfate, and concentrated, and then purified by column chromatograph (silica gel $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 100: 5) to give the product (2.75g).

5


TABLE 17

Ex. No.	R ₈	R ₉	HPLC retention time (min.)	MH ⁺
301	H	<i>t</i> -butyl	3.19	552
302	Et	Me	3.22	496


Example 303

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinepropanoic acid

10



Example 301 (200mg) was dissolved in 10 ml of ethanol. 1 ml of 10% LiOH/H₂O aqueous solution was added. The reaction solution was stirred at 45°C for 2 hours. Ethanol was removed under reduced pressure. The resulting aqueous solution was adjusted to pH=4 with 10% HCl. The solid was collected by filtration, rinsed with water, dried to afford 160mg slightly yellow solid. This product (300mg) was mixed in 0.3 ml of DMF and stirred at 130°C for 10 minutes. The reaction mixture was cooled to room temperature. Water was added. The solid was collected by filtration, rinsed with water, and dried to afford 244mg of the title compound. LC: 2.20', MH⁺: 396

Examples 304-3061-[(3-Chloro-4-methoxyphenyl)methyl]-1,2,3,4-tetrahydro-2-oxobenzo[h]-1,6-naphthyridine-9-carbonitriles

5

Compounds having the formula (Ias) wherein R_8 and X_2 have the values listed in Table 18, were prepared as follows.

- 10 Examples 304-305: To a suspension of NaH (123 mg) in THF (10 mL) was added dimethylmalonate (370 mg) dropwise. The mixture was stirred at RT for 15 min. then cooled to -78°C . A solution of 4-(3-chloro-4-methoxyphenylmethylamino)-3-chloromethyl-6-cyanoquinoline (0.85 mmol) in DMF was then added dropwise. The reaction was quenched at low
- 15 temperature and diluted with Et_2O then washed with water. The ether layer was concentrated and the residue was purified by flash column (silica gel, $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 100:4) to give example 303 (160 mg).

- Example 306: A mixture of the compound of Example 305 (20 mg), 1-
- 20 isopropylpyrrolizine (13 mg), EDAC•HCl (20 mg), HOBT• H_2O (15 mg) and DMAP (trace) in pyridine (1.5 mL) was stirred at RT for over night. The crude product was purified by preparative HPLC to give Example 306.

TABLE 18


Ex. No.	R_8	$-X_2$	HPLC	MP ⁺
---------	-------	--------	------	-----------------

			retention time (min.)	
304	H	-(C=O)OCH ₃	2.91	436
305	Et	-(C=O)OCH ₃	4.04	464
306	H	H	2.79	378

Examples 307-314

4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-N-alkyl-3-quinolinepropanamides

5









Compounds of formula (Iat) wherein R₁₀ and R₁₁ have the values listed in Table 19, were prepared as follows.

10 The compound of Example 303 (20mg, 0.05mmole) was mixed with BOP reagent (44 mg, 0.1mmole) in 3ml of pyridine and stirred at RT for 3 hours. The resulting reaction mixture was purified by preparative HPLC to afford a cream-colored solid.

15


TABLE 19

Ex. No.	-NR ₁₀ R ₁₁	HPLC retention time (min.)	MH ⁺
307		2.16	486
308	NH ₃	2.08	395

309		2.22	478
310		2.34	423
311		2.14	506
312		2.66	463
313		2.91	477
314		2.71	463

Example 3156-Bromo-N-[(3-chloro-4-methoxyphenyl)methyl]-2-(2-pyridinyl)-4-quinolinamine

5




To a suspension of 6-bromo-2-(2-pyridine)quinoline-3-aminobutyl ester (preparation 28) (300 mg) in DMF (10 mL) was added a solution of HMDSNa in THF (1 M, 0.9 mL) at RT to generate a brown solution. Then 3-chloro-4-methoxybenzyl chloride (160 mg) was added. The reaction mixture was stirred at RT for 2 h. An additional solution of HMDSNa in THF (0.9 mL) and 3-chloro-4-methoxybenzyl chloride (2 eq) was added and continue to stir. When the starting material was consumed, the reaction mixture was treated with TFA. Removal of the solvent and the residue was redissolved in EtOAc and washed with 1 N NaOH and water. Removal of the solvent and triturated

with Et₂O to give the title compound (163 mg) as a light yellow solid. LC:
1.73' MH⁺: 454.

What is claimed is:

1. A compound of the formula (I):



5

or a pharmaceutically-acceptable salt thereof, wherein:

R_2 , R_3 , R_7 and R_8 are independently hydrogen, halogen, alkyl, substituted alkyl, alkoxy, nitro, cyano, aryl, heteroaryl, or heterocyclo;

- 10 R_3 is $-(CH_2)_zY$, wherein z is 0, 1, 2, or 3;

R_4 and R_5 (i) are independently hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, or heteroaryl, with the proviso that R_4 and R_5 are not both hydrogen; (ii) taken together form a heterocyclo ring; or (iii) one of R_4 and R_5 together with Y forms a heterocyclo ring;

- 15 Y is (i) independently selected from $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$, $-O(C=O)NR_{10}R_{11}$, $-NR_{10}R_{11}$, $-NR_{10}(C=O)NR_{11}R_{12}$, $-CH[(C=O)NR_{10}R_{11}]_2$, $-(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, $-S(O)_mR_9$, $-SO_2NR_{10}R_{11}$, imidazole, substituted imidazole, triazole, substituted triazole, or cyano, or (ii) together with R_4 or R_5 forms a heterocyclo ring; and

- 20 m is 0, 1, or 2;

R_9 is hydrogen, alkyl, substituted alkyl, hydroxy, alkoxy, cycloalkyl, substituted cycloalkyl, heterocyclo, aryl, heteroaryl, or pentafluorophenyl; and

R_{10} , R_{11} , and R_{12} are (i) independently selected from hydrogen, alkyl, substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl, heterocyclo, and heteroaryl; or (ii) taken together, wherein R_{10} forms a three- to seven-membered heterocyclo ring with R_{11} or R_{12} , or R_{11} forms a three- to seven-membered heterocyclo ring with R_{12} .

25

2. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof, wherein:
- R_2 is hydrogen, halogen, lower alkyl, or heteroaryl;
- R_4 is hydrogen, lower alkyl, or forms a heterocyclo ring with Y;
- 5 R_5 is substituted alkyl having at least one substituent that is aryl, cycloalkyl, or heteroaryl;
- R_6 is hydrogen, halogen, trifluoromethyl, or cyano;
- R_7 is hydrogen or trifluoromethyl;
- R_8 is hydrogen, lower alkyl, or substituted alkyl having at least one substituent
- 10 that is $-(CH_2)_nNR_{13}, R_{14}$, wherein R_{13} and R_{14} (i) are independently selected from hydrogen, alkyl, and substituted alkyl, or (ii) taken together form a heterocyclo ring;
- n is 0, 1, 2, or 3;
- Y is (i) $-OR_9, -CO_2R_9, -CH(CO_2R_9)_2, -OR_9NR_{10}R_{11}, -NR_{10}R_{11},$
- 15 $-(C=O)NR_{10}R_{11}$, or $-NR_{10}(C=O)R_{12}$, or (ii) together with R_4 forms a heterocyclo ring;
- R_9 is hydrogen, alkyl, substituted alkyl, alkoxy, heterocyclo, or pentafluorophenyl;
- R_{10} and R_{11} are (i) independently selected from hydrogen, alkyl, substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl, heterocyclo, and
- 20 heteroaryl; or (ii) taken together wherein R_{10} and R_{11} forms a three- to seven-membered heterocyclo ring; and
- R_{12} is aryl, cycloalkyl, or heteroaryl.
- 25
3. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof, wherein:
- R_2 is hydrogen or chloro;
- R_4 is hydrogen;
- 30 R_5 is 3-chloro-4-methoxyphenylmethyl;
- R_6 is cyano;

R₇ is hydrogen; and

R₈ is hydrogen, alkyl or substituted alkyl.

4. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof,
5 wherein R₄ is hydrogen or lower alkyl, and R₅ is an alkyl substituted with
cycloalkyl or aryl.

5. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof,
in which R₅ is a substituted alkyl having a mono- or di-substituted aryl group
10 in which at least one of the aryl substituents is selected from halogen and
methoxy.

6. A compound of Claim 5 or a pharmaceutically-acceptable salt thereof,
wherein R₅ is 3-chloro-4-methoxyphenylmethyl.

15

7. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof,
wherein Y is (i) -OR₉, -CO₂R₉, -CH(CO₂R₉)₂, -OR₉NR₁₀R₁₁, -NR₁₀R₁₁, -
(C=O)NR₁₀R₁₁, or -NR₁₀(C=O)R₁₂, or (ii) together with R₄ forms a heterocyclo
ring.

20

8. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof,
wherein Y is CO₂R₉, and R₉ is hydrogen, lower alkyl, alkoxy, or substituted
alkyl having at least one substituent that is a heterocyclo group, wherein
when said heterocyclo group has the heteroatom N, said heteroatom N is
25 substituted with a group X₁ selected from alkyl, substituted alkyl, and
cycloalkyl.

9. A compound of Claim 1 or a pharmaceutically-acceptable salt thereof,
wherein R₃ is (CH₂)₂NR₁₀R₁₁, (C=O)NR₁₀R₁₁ or (CH₂)₂NR₉R₁₀R₁₁;
30 R₉ is alkoxy having from 1 to 4 carbon atoms, said alkoxy having a
substituent selected from hydroxy and alkoxy; and

R₁₀ and R₁₁ are (i) independently hydrogen, alkyl, substituted alkyl, cycloalkyl, alkoxy, or heterocyclo, or (ii) together form a heterocylco ring.

10. The compound of claim 1, wherein z is zero, R₃ is OR₉, and R₉ is
5 hydrogen or lower alkyl.


11. The compound of claim 1, wherein R₆ is cyano.

12. The compound of claim 1 selected from

- 10 (i) 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-8-ethyl-N-(2-pyridinylmethyl)-3-quinolinecarboxamide; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-(hydroxymethyl)-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-[(3R)-3-hydroxy-1-pyrrolidinyl]methyl]-6-quinolinecarbonitrile; N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinyl]methyl]benzamide; N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinyl]methyl]cyclohexanecarboxamide; N-[[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinolinyl]methyl]-2-pyridinecarboxamide; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-(propoxymethyl)-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-(methoxymethyl)-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[[[tetrahydro-2-furanyl)methoxy]methyl]-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[(1-methylethoxy)methyl]-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-(ethoxymethyl)-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-[(2-ethoxyethoxy)methyl]-8-ethyl-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[(2-hydroxyethoxy)methyl]-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[(oxiranylmethoxy)methyl]-6-
- 15
- 20
- 25
- 30

- quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[[[(2S)-1-methyl-2-pyrrolidinyl]methoxy]methyl]-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[[[(2S)-1-cyclopentyl-2-pyrrolidinyl]methoxy]methyl]-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-8-ethyl-3-[[2-hydroxy-3-(1-pyrrolidinyl)propoxy]methyl]-6-quinolinecarbonitrile; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-3-[[3-(dimethylamino)-2-hydroxypropoxy]methyl]-8-ethyl-6-quinolinecarbonitrile; [[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-8-ethyl-3-quinoliny]]methyl]propanedioic acid dimethyl ester; 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-N-ethyl-3-quinolinepropanamide; 1-[3-[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinoliny]]-1-oxopropyl]-4-(1-methylethyl)piperazine; 1-[3-[4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-3-quinoliny]]-1-oxopropyl]piperidine, trifluoroacetate; or 4-[[[(3-Chloro-4-methoxyphenyl)methyl]amino]-6-cyano-N-cyclopentyl-3-quinolinepropanamide; and (ii) a pharmaceutically-acceptable salt thereof.

- 20 13. A pharmaceutical composition adapted for treating a cGMP-associated condition comprising (a) one or more compounds of the formula (I),



- 25 or a pharmaceutically acceptable salt thereof, and (b) a pharmaceutically acceptable carrier or diluent, wherein:
R₂, R₆, R₇ and R₈ are independently hydrogen, halogen, alkyl, substituted alkyl, alkoxy, nitro, cyano, aryl, or heteroaryl;

- R_3 is hydrogen or $-(CH_2)_zY$, wherein z is 0, 1, 2, or 3;
- R_4 and R_5 (i) are independently hydrogen, alkyl, substituted alkyl, cycloalkyl, substituted cycloalkyl, aryl, or heteroaryl, with the proviso that R_4 and R_5 are not both hydrogen; or (ii) taken together form a heterocyclo ring;
- 5 or (iii) one of R_4 and R_5 together with Y forms a heterocyclo ring;
- Y is selected (i) independently from $-OR_9$, $-CO_2R_9$, $-CH(CO_2R_9)_2$, $-O(C=O)NR_{10}R_{11}$, $-NR_{10}R_{11}$, $-NR_{10}(C=O)NR_{11}R_{12}$, $-CH[(C=O)NR_{10}R_{11}]_2$, $-(C=O)NR_{10}R_{11}$, $-NR_{10}(C=O)R_{12}$, $-S(O)_mR_9$, $-SO_2NR_{10}R_{11}$, imidazole, substituted imidazole, triazole, substituted triazole, and cyano, or (ii)
- 10 together with R_4 or R_5 , forming a heterocyclo ring therewith;
- m is 0, 1, or 2;
- R_9 is hydrogen, alkyl, substituted alkyl, hydroxy, alkoxy, cycloalkyl, substituted cycloalkyl, heterocyclo, aryl, heteroaryl, or pentafluorophenyl; and
- R_{10} , R_{11} , and R_{12} (i) are independently selected from hydrogen, alkyl,
- 15 substituted alkyl, alkoxy, cycloalkyl, substituted cycloalkyl, aryl, heterocyclo, and heteroaryl; or (ii) taken together wherein R_{10} forms a three- to seven-membered heterocyclo ring with R_{11} or R_{12} , or R_{11} forms a three- to seven-membered heterocyclo ring with R_{12} .
- 20 14. The pharmaceutical composition of claim 13, wherein:
- R_2 is hydrogen or chloro;
- R_4 is hydrogen;
- R_5 is 3-chloro-4-methoxyphenylmethyl;
- R_6 is cyano;
- 25 R_7 is hydrogen; and
- R_8 is hydrogen, alkyl or substituted alkyl.

15. A method of treating a cGMP-associated condition comprising
- 30 administering to a mammal in need thereof a therapeutically-effective amount of one or more compounds of Claim 1.

16. A method of treating a cGMP-associated condition comprising administering to a mammal in need thereof a therapeutically-effective amount of a composition according to Claim 13.
- 5
17. A method of treating a cardiovascular disorder comprising administering to a mammal in need thereof a therapeutically-effective amount of a composition according to Claim 13.
- 10 18. A method of treating sexual dysfunction in a male or female mammal comprising administering to a mammal in need thereof a therapeutically-effective amount of a composition according to Claim 13.
- 15 19. A method of treating diabetes comprising diabetes mellitus comprising administering to a mammal in need thereof a therapeutically-effective amount of a composition according to Claim 13.
- 20 20. A method of treating gastrointestinal disorders comprising gastric paresis comprising administering to a mammal in need thereof a therapeutically-effective amount of a composition according to Claim 13.